

Booster Dose

A single Janssen COVID-19 Vaccine booster dose (0.5 mL) may be administered at least 2 months after primary vaccination with the Janssen COVID-19 Vaccine, to individuals 18 years of age and older.

A single booster dose of the Janssen COVID-19 Vaccine (0.5 mL) may be administered to individuals 18 years of age and older as a heterologous booster dose following completion of primary vaccination with another authorized or approved COVID-19 vaccine. The dosing interval for the heterologous booster dose is the same as that authorized for a booster dose of the vaccine used for primary vaccination.

3 DOSAGE FORMS AND STRENGTHS

Janssen COVID-19 Vaccine is a suspension for intramuscular injection. A single-dose is 0.5 mL.

4 CONTRAINDICATIONS

4.1 Severe Allergic Reactions

Do not administer the Janssen COVID-19 Vaccine to individuals with a known history of severe allergic reaction (e.g., anaphylaxis) to any component of the Janssen COVID-19 Vaccine [*see Description (13)*].

4.2 Thrombosis with Thrombocytopenia

Do not administer the Janssen COVID-19 Vaccine to individuals with a history of thrombosis with thrombocytopenia following the Janssen COVID-19 Vaccine or any other adenovirus-vectored COVID-19 vaccine (e.g., AstraZeneca's COVID-19 vaccine which is not authorized or approved in the United States) [*see Warnings and Precautions (5.2)*].

5 WARNINGS AND PRECAUTIONS

5.1 Management of Acute Allergic Reactions

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of the Janssen COVID-19 Vaccine.


Monitor Janssen COVID-19 Vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

5.2 Thrombosis with Thrombocytopenia Syndrome (TTS)

Reports to the Vaccine Adverse Events Reporting System (VAERS), a passive surveillance system, provide evidence for an increased risk of thrombosis with thrombocytopenia syndrome (TTS) with onset of symptoms approximately one to two weeks after administration of the Janssen COVID-19 Vaccine.

21 CONTACT INFORMATION

For general questions or to access the most recent Janssen COVID-19 Vaccine Fact Sheets, scan the QR code using your device, visit www.janssencovid19vaccine.com or call the telephone numbers provided below.

QR Code	Fact Sheets Website	Telephone numbers
	www.janssencovid19vaccine.com	US Toll Free: 1-800-565-4008 US Toll: 1-908-455-9922

This Full EUA Prescribing Information may have been updated. For the most recent Full EUA Prescribing Information, please see www.janssencovid19vaccine.com.

Manufactured by:
Janssen Biotech, Inc.
a Janssen Pharmaceutical Company of Johnson & Johnson
Horsham, PA 19044, USA



Revised: JAN/31/2022
© 2021 Janssen Pharmaceutical Companies