

Full Reference List for BPA Draft Assessment

References are only listed once in order of their first appearance in the assessment.

- 1.) FDA's Preparation of Food Contact Notifications for Food Contact Substances: Toxicology Recommendations Final Guidance April 2002 (accessible at <http://www.cfsan.fda.gov/~dms/opa2pmnt.html>).
- 2.) Krishnan, AV, Stathis P, Permuth SF, Tokes L, and Feldman D. (1993) Bisphenol A: An estrogenic substance is released from polycarbonate flasks during autoclaving. *Endocrinology* 132(6): 2279-2286.
- 3.) Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, Van Der Saag PT, Van Der Burg B, and Gustafsson J-A. (1998) Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinology* 139(10) 4252-4263.
- 4.) Summarized data cited in CERHR final report NTP-CERHR Expert Panel Report on the Reproductive and Developmental Toxicity of Bisphenol A, dated November 2007 (accessible at <http://cerhr.niehs.nih.gov/chemicals/bisphenol/BPAFinalEPVF112607.pdf>) and published as Chapin et al. (2008) *NTP-CERHR Expert Panel Report on the Reproductive and Developmental Toxicity of Bisphenol A Birth Defects Research* (Part B) 83:157-395.
- 5.) Melnick R, Lucier G, Wolfe M, Hall R, Stancel G, Prins G, Gallo M, Reuhl K, Ho SM, Brown T, Moore J, Leakey J, Haseman J, Kohn M (2002) *Environ Health Perspect*. Summary of the National Toxicology Program's report of the endocrine disruptors low-dose peer review. 110(4): 427-431.
- 6.) Jensen, MN and Ritskes-Hoitinga, M. (2007) How isoflavone levels in common rodent diets can interfere with the value of animal models and with experimental results *Lab Anim* 41(1):1-18.
- 7.) Brown, NM and Setchell, KD. (2001) Animal models impacted by phytoestrogens in commercial chow: implications for pathways influenced by hormones. *Lab Invest* 81(5):735-47.
- 8.) Thigpen JE, Setchell KD, Ahlmark KB, Locklear J, Spahr T, Caviness GF, Goelz MF, Haseman JK, Newbold RR, Forsythe DB (1999) Phytoestrogen content of purified, open- and closed-formula laboratory animal diets. *Lab Anim Sci* 49(5):530-6.
- 9.) Setchell, KD (1998) Phytoestrogens: the biochemistry, physiology, and implications for human health of soy isoflavones *Am J Clin Nutr* 68: 1333S-1346S
- 10.) Mazur, W and Adlercreutz, H. (2000) Overview of Naturally Occurring Endocrine-Active Substances in the Human Diet in Relation to Human Health *Nutrition* 16:654-687
- 11.) Cassidy, A and Setchell, KDR (1999) Dietary Isoflavones: Biological Effects and Relevance to Human Health *J Nutr* 129(3):758S-767S.
- 12.) Bhatia J, Greer, F and the Committee on Nutrition (2008) Use of Soy Protein-Based Formulas in Infant Feeding *Pediatrics* 121;1062-1068
- 13.) Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids and Materials in Contact with Food on a request from the Commission related to 2,2-Bis(4-Hydroxyphenyl)Propane

(Bisphenol A) Question number EFSA-Q-2005-100 Adopted on 29 November 2006, *The EFSA Journal* (2006) 428, 1 – 75.

14.) Bisphenol A Risk Assessment Document, November 2005, (AIST Risk Assessment Document Series No. 6) New Energy and Industrial Technology Development Organization (NEDO) and Research Center for Chemical Risk Management (CRM) Japanese National Institute of Advanced Industrial Science and Technology (AIST) original 2005 document; a 2007 document is currently available.

15.) Draft NTP Brief on Bisphenol A, April 14th, 2008. Accessible at http://cerhr.niehs.nih.gov/chemicals/bisphenol/BPADraftBriefVF_04_14_08.pdf

16.) Government of Canada, Environment Canada draft screening assessment and risk management documents dated April 2008 accessible at http://www.chemicalsubstanceschimiques.gc.ca/challenge-defi/batch-lot_2_e.html#ReleaseofDraft

17.) Updated European Risk Assessment Report 4,4'-Isopropylidenediphenol (Bisphenol-A) CAS Number: 80-05-7 EINECS Number: 201-245-8 Final Approved Version Awaiting For Publication accessible at http://ecb.jrc.it/documents/Existing-Chemicals/RISK_ASSESSMENT/ADDENDUM/bisphenola_add_325.pdf

18.) Opinion of the Scientific Panel on Food Additives, Flavourings, Processing Aids, Materials in Contact with Food and Cosmetics of the Norwegian Scientific Committee for Food Safety 18 June 2008 Assessment of four studies on developmental neurotoxicity of bisphenol A, accessible at http://www.vkm.no/eway/default.aspx?pid=266&trg=MainLeft_5419&MainLeft_5419=5468:17924::0:5420:1::0:0

19.) European Food Safety Authority, Toxicokinetics of Bisphenol A Scientific Opinion of the Panel on Food additives, Flavourings, Processing aids and Materials in Contact with Food (AFC) (Question No EFSA-Q-2008-382) Adopted on 9 July 2008 http://www.efsa.eu.int/EFSA/efsa_locale-1178620753812_1211902017492.htm

20.) Full text of the 2006 opinion of the former Panel on Food Additives, Flavourings, Processing aids and Materials in Contact with Food (AFC) at http://www.efsa.eu.int/EFSA/efsa_locale-1178620753812_1178620772817.htm

21.) Environmental Working Group. Bisphenol A: Toxic Plastics Chemical in Canned Food: Canned food exposures are significant, accessible at <http://www.ewg.org/node/20936> and Toxic Plastics Chemical in Infant Formula, accessible at <http://www.ewg.org/reports/bpaformula>.

22.) *Reproductive Toxicology* 24(2):August-September 2007

23.) Integrated Risk Information System (IRIS), last update 1993 accessible <http://www.epa.gov/ncea/iris/subst/0356.htm>

24.) U.S. Environmental Protection Agency mode of action cross-species informational assessment accessible at <http://cfpub.epa.gov/ncea/cfm/recorddisplay.cfm?deid=22445>

25.) Effective FCNs include 362, 363, 404, 463, 517, 624, 702, and 737; details are available at <http://www.cfsan.fda.gov/~dms/opa-fcn.html>

- 26.) FDA Memorandum – Memorandum to the File. FAP 9Z4681 (MATS #1070 M2.0 and 2.1): National Environmental Trust; submission of 5/13/99. Migration of di(2-ethylhexyl) adipate from polyvinyl chloride cling film and bisphenol-A from can coatings and polycarbonate baby bottles and tableware. Paquette/Smith, 05/2/2000
- 27.) FDA Memorandum – Memorandum to the File. Cumulative exposure estimates for bisphenol A (BPA). Bailey/Cheeseman, 08/13/2001.
- 28.) FDA Memorandum – Memorandum to the File. Cumulative exposure estimates for bisphenol A (BPA), individually for adults and infants, from its use in epoxy-based can coatings and polycarbonate (PC) articles. Verbal request of 10-23-95. Bailey/Diachenko, 03/13/2006.
- 29.) FDA Memorandum – Memorandum to the File. A summary of FDA's Chemistry Methods Branch's (CMB) studies on BPA migration from PC infant bottles and BPA level in vegetables was provided to the Chemistry Review Branch (CRB) on 9-26-95 by CMB (Henry Hollifield) in a draft report entitled "Bisphenol-A: Status Summary Report."
- 30.) FDA Memorandum – Memorandum to the File. CMB's studies on BPA levels in infant formula are described in a CMB memorandum dated 3-5-96 (J. Biles to G. Diachenko).
- 31.) Biles, J., McNeal, T., Begley, T., Hollifield, H. (1997) Determination of Bisphenol-A in Reusable Polycarbonate Food-Contact Plastics and Migration to Food-Simulating Liquids., *Journal of Agricultural and Food Chemistry*, Vol. 45(9): 3541-3544.
- 32.) Biles, J., McNeal, T., Begley. (1997) Determination of Bisphenol-A Migrating from Epoxy Can Coatings to Liquid Infant Formula Concentrates. *Journal of Agricultural and Food Chemistry* 45(9): 4697-4700.
- 33.) Brotons, J.A., Olea-Serrano, M.F., Villalobos, M., Pedraza, V., and Olea, N. (1995) Xenoestrogens Released from Lacquer Coatings in Food Cans, *Environmental Health Perspectives* 103:609-612.
- 34.) Guidance for Industry Preparation of Premarket Submissions for Food Contact Substances: Chemistry Recommendations (accessible at <http://www.cfsan.fda.gov/~dms/opa3pmnc.html>);
- 35.) FDA Memorandum - Memorandum to the File. Update on cumulative exposure to BPA for infants from epoxy-based container coatings and polycarbonate (PC) bottles in contact with infant formula. Verbal request dated 4/29/08. Bailey/Twaroski, 06/02/2008.
- 36.) USDA Continuing Survey of Food Intakes by Individuals data accessible at http://www.ars.usda.gov/main/site_main.htm?modecode=12-35-50-00
- 37.) National Health and Nutrition Examination Survey (NHANES), 2003-2004, accessible at http://www.cdc.gov/nchs/about/major/nhanes/nhanes2003-2004/exam03_04.htm.
- 38.) FDA Memorandum – Memorandum to the File. Results from W. Limm (HFS-706) on two brands of infant formula packaged in composite containers indicates that the linings not based on epoxy chemistry. Memorandum dated 5/28/08, W. Limm to A. Bailey.
- 39.) Infant body masses taken from CDC (accessible at <http://www.cdc.gov/nchs/about/major/nhanes/growthcharts/datafiles.htm>)

- 40.) Calafat, A.M., Ye, X., Wong, Y-L., Reidy, J.A. and Needham and L.L. (2008) Exposure of the U.S. Population to Bisphenol A and 4-tertiary-Octylphenol:2003–2004 *Environ Health Perspect* 116:39–44
- 41.) Goodman, J.E., McConnell, E. E, Sipes, I.G., Witorsch, R.J., Slayton, T.M., Yu, C.J., Lewis, A.S., Rhombert, L.R. (2006) An Updated Weight of the Evidence Evaluation of Reproductive and Developmental Effects of Low Doses of Bisphenol A. *Critical Reviews in Toxicology*, 36:387-457
- 42.) Haighton, L.A., Hlywka, J.J., Doull, J., Kroes, R. Lynch, B.S., and Munro, I.C. (2002) An Evaluation of the Possible Carcinogenicity of Bisphenol A to Humans, *Regulatory Toxicology and Pharmacology* 35:238-254
- 43.) Willhite, C.C., Ball. G.L. and McLellan, C.J. (2008). Derivation of a bisphenol A oral reference dose (RfD) and drinking-water equivalent concentration. *Journal of Toxicology and Environmental Health, Part B*, 11:2, 69 – 146
- 44.) Gray GM, Cohen JT, Cunha G, Hughes C, McConnell EE, Rhomberg L, Sipes IG, Mattison D (2004) Weight of the evidence evaluation of low-dose reproductive and developmental effects of bisphenol A. *Human and Ecological Risk Assessment* 10(5): 875-921
- 45.) Huff, J. (2001) Carcinogenicity of Bisphenol A in Fischer 344 and B6CC3F1 mice. *Odontology* 89:12-20.
- 46.) The five levels of concern used by NTP are from highest to lowest: serious concern, concern, some concern, minimal concern, and negligible concern. Definitions of these levels are not defined by NTP. <http://www.niehs.nih.gov/news/media/questions/sya-bpa.cfm>
- 47.) Actions on the Draft NTP Brief on Bisphenol A by the NTP Board of Scientific Counselors (BSC), June 11, 2008, accessible at http://ntp.niehs.nih.gov/files/BSCactionsBPA_508.pdf.
- 48.) Guidance for Industry and Other Stakeholders Toxicological Principles for the Safety Assessment of Food Ingredients Redbook 2000 accessible at <http://www.cfsan.fda.gov/~redbook/red-toca.html>
- 49.) Good Laboratory Practices, 21 CFR §58.1 (Title 21 of the Code of Federal Regulations, Section 58.1), accessible at http://edocket.access.gpo.gov/cfr_2008/aprqrtr/21cfr58.1.htm
- 50.) Matthews, J.B., Twomey, K., Zacharewski, T.R. (2001) *In vitro* and *in vivo* interactions of bisphenol A and its metabolite, bisphenol A glucuronide, with estrogen receptors alpha and beta. *Chem Res Toxicol.* 2:149-57.
- 51.) FDA Memorandum - Compact Summary of Bisphenol A (BPA) Pharmacokinetics. Roth and Komolprasert/Twaroski, 06/01/2007; revised 05/23/2008.
- 52.) Ye, X., Bishop, L., and Needham, L., and Calafat A.M. (2008) Automated on-line column-switching HPLC-MS/MS method with peak focusing for measuring parabens, triclosan, and other environmental phenols in human milk *Anal Chem Acta* 622(1-2):150-6.
- 53.) Ye, X, Kuklenyik, Z., Needham, L.L. and Calafat., A.M. (2006) Measuring environmental phenols and chlorinated organic chemicals in breast milk using automated on-line column-switching–high performance liquid chromatography–isotope dilution tandem mass spectrometry. *J. Chromatography B* 2006;831:110-115.

- 54.) Carcinogenesis Bioassay of Bisphenol A in F344 rats and B6C3F1 mice - Feed Study, NTP Technical Report 215, NIH Publication No. 82-1761.
- 55.) FDA Review Memorandum -Acceptance of Final TDERs for review of NTP's Carcinogenesis Bioassay of Bisphenol A in F344 rats and B6C3F1 mice (Feed Study) (NTP TR 215). Shackelford/Food Additive Master file 580. 07/24/2007
- 56.) EU Risk Assessment Report on BPA – Final Report, 2003 accessible at http://ecb.jrc.it/DOCUMENTS/Existing-Chemicals/RISK_ASSESSMENT/REPORT/bisphenolareport325.pdf
- 57.) FDA Memorandum - CAC Meeting Dates: 04/24/2008, 05/09/2008 CFSAN Cancer Assessment Committee (CAC), Full CAC Review – Bisphenol A (BPA)
- 58.) FDA Review Memorandum - Acceptance of Final TDERs for studies reviewed under contract with ICF Consulting (Contract Number 223-96-2302) for Food Additive Master File No 580 under Work Assignment 2000-20 (ICF 020) Tasks Number 1, 2 and 3. Shackelford/Food Additive Master File 580, 07/24/2007.
- 59.) FDA memorandum Shackelford/Twaroski, 06/24/2007: Review of Two-Generation Reproductive Toxicity Evaluation of Bisphenol A Administered in the Feed to CD-1® Swiss Mice; RTI study number (Study number 65C-09301.000.003/0209301.000.003) published as Tyl RW, Myers CB, Marr MC, Sloan CS, Castillo NP, Veselica MM, Seely JC, Dimond SS, Van Miller JP, Shiotsuka RN, Beyer D, Hentges SG, Waechter JM Jr (2008) Two-generation reproductive toxicity study of dietary bisphenol A (BPA) in CD-1 (Swiss) mice. *Toxicol Sci* 104(2): 362-384.
- 60.) FDA memorandum Gu/Twaroski, 07/18/2007: Review of study entitled “Three-Generation Reproductive Toxicity Evaluation of Bisphenol A in the Feed of CD® (Sprague-Dawley) Rats” RTI study number 65C-07036-000 published as Tyl RW, Myers CB, Marr MC, Thomas BF, Keimowitz AR, Brine DR, Veselica MM, Fail PA, Chang TY, Seely JC, Joiner RL, Butala JH, Dimond SS, Cagen SZ, Shiotsuka RN, Stropp GD, Waechter JM (2002) *Toxicol Sci*. Three-generation reproductive toxicity study of dietary bisphenol A in CD Sprague-Dawley rats. 68(1): 121-146.
- 61.) FDA e-mail “About AGD” dated 06/06/2008 (Gu/Twaroski).
- 62.) Ema, M., Fujii, S., Furukawa, M., Kiguchi, M., Ikka, T., and Harazono, A. (2001): Rat two-generation reproductive toxicity study of bisphenol A. *Reprod Toxicol* 15:505-523
- 63.) FDA Memorandum - Table of NOAELS and LOAELS from Bisphenol A Toxicity Studies in FMF 580. Shackelford/Twaroski, 06/12/2007.
- 64.) FDA Memorandum - Acceptance of Final TDERs for studies reviewed under contract with ICF Consulting (contract No. 223-96-2302) for Food Additive Master File No. 580 under Work Assignment 2000-19 (ICF 419), Task Numbers 3, 4, 5, 6. Shackelford/Twaroski, 06/11/2007.
- 65.) FDA Memorandum – Acceptance of “Bisphenol A – Effects on onset of puberty in female and prostate and urinary tract in male rodents” reviewed by Drs. K. Barry Delclos (HFT-110) and Deborah K. Hansen (HFT-130) at FDA's National Center for Toxicological Research (NCTR). Twaroski/Gu/Food Master File 580. 05/27/2008.
- 66.) Safranski ,T.J., Lamberson, W.R. and Keisler, D.H. (1993) Correlations among three measures of

puberty in mice and relationships with estradiol concentration and ovulation *Biology of Reproduction*, 48: 669-673.

67.) Nelson, JF, Karelus, K, Felicio, LS and Johnson TE (1990) Genetic influences on the timing of puberty in mice. *Biol of Reproduction* 42: 649-655.

68.) Cooper RL et al. In: Heindel, J.J., Chapin, R.E. (Eds.), Female Reproductive Toxicology. Methods Toxicol., Vol. 3B. Academic Press, pp. 45–56.

69.) Written comments provided by Gray LE (former member of the CERHR BPA Expert Panel) on the NTP draft brief (05/23/2008, accessible at [http://cerhr.niehs.nih.gov/chemicals/bisphenol/pubcomm/BPA\(37\)Gray23May2008best.pdf](http://cerhr.niehs.nih.gov/chemicals/bisphenol/pubcomm/BPA(37)Gray23May2008best.pdf).

70.) Euling, S.Y., Selevan, S.G., Pescovitz, O.H. and Skakkebaek, N.E. (2008) Role of Environmental Factors in the Timing of Puberty. *Pediatrics* 121;S167-S171.

71.) FDA Memorandum – Sprando/Biddle, Review of MPI report on bisphenol A, 02/04/1999, cover memo (Twaroski/Food Master File 580, 01/22/2008), published as Cagen, S.Z., Waechter, J.M., Dimond, S.S., Breslin, W.J., Butala, J.H., Jekat, F.W., Joiner, R.L., Shiotsuka, R.N., Veenstra, G.E., and Harris, L.R. (1999) Normal reproductive organ development in CF-1 mice following prenatal exposure to bisphenol A, *Toxicological Sciences*, 50: 36-44.

72.) FDA Memorandum - Acceptance of updated reviews of the developmental neurotoxicity potential performed by Oak Ridge National Laboratory (ORNL, FDA Interagency Agreement #224-00-2615, Task #2007-20) and by Drs. Sherry A. Ferguson and Merle G. Paule at FDA's National Center for Toxicological Research Food Master File 580. 05/28/2008.

73.) Wilson, C.A. and Davies, D.C. (2007) The control of sexual differentiation of the reproductive system and brain. *Reproduction* 133:331-359.

74.) Collins TFX, Sprando RL, Shackelford ME, Gruber MF, and Morse DE. (2006) Principles of Risk Assessment – FDA Perspective. In Developmental and Reproductive Toxicology- A Practical Approach, Taylor & Francis (Ed. Ronald D. Hood). p. 877-909.

75.) Twaroski, M.L., Batarseh, L.I., and Bailey, A.B. The Regulation of Food Contact Substances in the United States. Chemical Migration and Food Contact Materials. Woodhead Publishing (2007) Barnes, Sinclair, and Watson (Eds.).

76.) Meeting Presentations – June 11 – 12, 2008 BSC, accessible at <http://ntp.niehs.nih.gov/index.cfm?objectid=78A617B9-F1F6-975E-7F3871DF6BC95C22>

77.) OECD Guideline For The Testing Of Chemicals Draft Proposal For A New Guideline 426 Developmental Neurotoxicity Study accessed at <http://www.oecd.org/dataoecd/20/52/37622194.pdf>

78.) Ho SM, Tang WY, Belmonte de Frausto J, Prins GS (2006) *Cancer Res*. Developmental exposure to estradiol and bisphenol A increases susceptibility to prostate carcinogenesis and epigenetically regulates phosphodiesterase type 4 variant 4. 66(11): 5624-5632.

79.) Timms BG, Howdeshell KL, Barton L, Bradley S, Richter CA, vom Saal FS (2005) *Proc Natl Acad Sci U S A*. Estrogenic chemicals in plastic and oral contraceptives disrupt development of the fetal mouse prostate and urethra. 102(19): 7014-7019.

- 80.) Ogura Y, Ishii K, Kanda H, Kanai M, Arima K, Wang Y, Sugimura Y (2007) *Differentiation*. Bisphenol A induces permanent squamous change in mouse prostatic epithelium. 75(8): 745-756.
- 81.) Durando M, L. K, Piva J, Sonnenschein C, Soto AM, Luque E, Muñoz-de-Toro M (2007) *Environ Health Perspect*. Prenatal bisphenol A exposure induces preneoplastic lesions in the mammary gland in Wistar rats. 11580-86.
- 82.) Murray TJ, Maffini MV, Ucci AA, Sonnenschein C, Soto AM (2007) *Reprod Toxicol*. Induction of mammary gland ductal hyperplasias and carcinoma in situ following fetal bisphenol A exposure. 23(3): 383-390.
- 83.) Markey CM, Luque EH, Munoz De Toro M, Sonnenschein C, Soto AM (2001) *Biol Reprod*. In utero exposure to bisphenol A alters the development and tissue organization of the mouse mammary gland. 65(4): 1215-1223 [Erratum: *Biol Reprod* 2004;1271:1753].
- 84.) Moral R, Wang R, Russo IH, Lamartiniere CA, Pereira J, Russo J (2008) *J Endocrinol*. Effect of prenatal exposure to the endocrine disruptor bisphenol A on mammary gland morphology and gene expression signature. 196(1): 101-112.
- 85.) Howdeshell KL, Hotchkiss AK, Thayer KA, Vandenberg JG, vom Saal FS (1999) *Nature*. Exposure to bisphenol A advances puberty. 401(6755): 763-764.
- 86.) Honma S, Suzuki A, Buchanan DL, Katsu Y, Watanabe H, Iguchi T (2002) *Reprod Toxicol*. Low dose effect of in utero exposure to bisphenol A and diethylstilbestrol on female mouse reproduction. 16(2): 117-122.
- 87.) Ryan BC, Vandenberg JG (2006) *Horm Behav*. Developmental exposure to environmental estrogens alters anxiety and spatial memory in female mice. 50(1): 85-93.
- 88.) Ashby J, Tinwell H, Haseman J (1999) *Regul Toxicol Pharmacol*. Lack of effects for low dose levels of bisphenol A and diethylstilbestrol on the prostate gland of CF1 mice exposed in utero. 30(2 Pt 1): 156-166.
- 89.) Markey CM, Coombs MA, Sonnenschein C, Soto AM (2003) *Evol Dev*. Mammalian development in a changing environment: exposure to endocrine disruptors reveals the developmental plasticity of steroid-hormone target organs. 5(1): 67-75.
- 90.) Negishi T, Kawasaki K, Suzaki S, Maeda H, Ishii Y, Kyuwa S, Kuroda Y, Yoshikawa Y (2004) *Environ Health Perspect*. Behavioral alterations in response to fear-provoking stimuli and tranlycypromine induced by perinatal exposure to bisphenol A and nonylphenol in male rats. 112(11): 1159-1164.
- 91.) Laviola G, Gioiosa L, Adriani W, Palanza P (2005) *Brain Res Bull*. D-amphetamine-related reinforcing effects are reduced in mice exposed prenatally to estrogenic endocrine disruptors. 65(3): 235-240.
- 92.) Gioiosa L, Fissore E, Ghirardelli G, Parmigiani S, Palanza P (2007) *Horm Behav*. Developmental exposure to low-dose estrogenic endocrine disruptors alters sex differences in exploration and emotional responses in mice. 52(3): 307-316.

- 93.) Ceccarelli I, Della Seta D, Fiorenzani P, Farabollini F, Aloisi AM (2007) *Neurotoxicol Teratol*. Estrogenic chemicals at puberty change ERalpha in the hypothalamus of male and female rats. 29(1): 108-115.
- 94.) Della Seta D, Minder I, Belloni V, Aloisi AM, Dessi-Fulgheri F, Farabollini F (2006) *Horm Behav*. Pubertal exposure to estrogenic chemicals affects behavior in juvenile and adult male rats. 50(2): 301-307.
- 95.) Palanza PL, Howdeshell KL, Parmigiani S, vom Saal FS (2002) *Environ Health Perspect*. Exposure to a low dose of bisphenol A during fetal life or in adulthood alters maternal behavior in mice. 110(Suppl 3): 415-422.
- 96.) Adriani W, Seta DD, Dessi-Fulgheri F, Farabollini F, Laviola G (2003) *Environ Health Perspect*. Altered profiles of spontaneous novelty seeking, impulsive behavior, and response to D-amphetamine in rats perinatally exposed to bisphenol A. 111(4): 395-401. Erratum (2005)
- 97.) Carr R, Bertasi F, Betancourt A, Bowers S, Gandy BS, Ryan P, Willard S (2003) *J Toxicol Environ Health A*. Effect of neonatal rat bisphenol A exposure on performance in the Morris water maze. 66(21): 2077-2088.
- 98.) "Baby Alert: New Findings About Plastics," *Consumer Reports*, May 1999, pp. 28-29.
- 99.) Takao, Y., Lee, H. C., Ishibashi, Y., Kohra, S., Tominaga, N., Arizono, K., "Fast Screening Method for Bisphenol A in Environmental Water and in Food by Solid-Phase Microextraction (SPME)," *Journal of Health Science*, Vol. 45, 1999, p. 39.
- 100.) Cited by NET as unpublished results by Iguchi of Yokohama City University, Japan.
- 101.) Simoneau, C., Roeder, G., Anklaam, E., "Migration of bisphenol-A from baby bottles: effect of experimental conditions and European survey." 2nd International Symposium on Food Packaging: Ensuring the Safety and Quality of Foods (ILSI conference), Vienna, Austria, 8-10 November 2000. (reference not available - cited from the EFSA report)
- 102.) Cited in the report as: Hanai, Y., "Bisphenol-A Eluted from Nursing Bottles," Unpublished Data, Environmental Science Research Center, Yokohama National University, 1997.
- 103.) Earls, A., Clay, C., Braybrook, J., 2000, "Preliminary Investigation into the Migration of Bisphenol-A from Commercially-Available Polycarbonate Baby Feeding Bottles". LGC Technical Report LGC/DTI/2000/005. (reference not available - cited from the EFSA report)
- 104.) Brede, C., Fjeldal, P., Skjevrak, I., Herikstad, H., "Increased migration levels of bisphenol A from polycarbonate baby bottles after dishwashing, boiling and brushing", *Food Additives and Contaminants*, Vol. 20, 2003, pp. 684-9.
- 105.) Tan, B., Mustafa, A. M., "Leaching of bisphenol A from new and old babies bottles, and new babies feeding teats", *Asia Pacific Journal of Public Health*, Vol. 15, 2003, pp. 118-23.
- 106.) A study of the migration of bisphenol A from polycarbonate feeding bottles into food simulants, Central Science Laboratory Test Report L6BB-1008 for the Boots Group, 2003 (available at www.boots-plc.com/environment/library/250.pdf).

- 107.) European Commission (2002), "Final opinion of the Scientific Committee on Food on Bisphenol A," April 17, 2002, SCF/CS/PM/3936 (http://ec.europa.eu/food/fs/sc/scf/out128_en.pdf).
- 108.) Kuo, H., Ding, W., "Trace determination of bisphenol A and phytoestrogens in infant formula powders by gas chromatography-mass spectrometry", *Journal Chromatography A*, Vol. 1027, 2004, pp. 67-74.
- 109.) A Survey of Bisphenol A in US Canned Foods, dated 3/5/07 (available on the Environmental Working Group website).
- 110.) Goodson, A., Summerfield, W., Cooper, I., "Survey of bisphenol A and bisphenol F in canned foods", *Food Additives and Contaminants*, Vol. 19, 2002, pp. 796-802.
- 111.) Personal communication between A. Bailey and T. Begley (HFS-706), May 2008.
- 112.) Lacroix, G., Lalonde, P., Feeley, M., "Bisphenol A (BPA) content of commercial polycarbonate (PC) baby bottles," Page, D., Health Canada Science Forum Book of Abstracts. 2006.
- 113.) Cao, X. Corriveau J., 2008, "Migration of bisphenol A from polycarbonate baby and water bottles into water under severe conditions", *J Agric. Food Chem.* 56(15): 6378-81 .
- 114.) Health Canada, Safe Environment Programme, HECSB, personal communication, 3/6/08, unreferenced.
- 115.) "Toxic baby bottles in Canada,": 2008, Environmental Defence (available at <http://www.toxicnation.ca/files/toxicnation/report/ToxicBabyBottleReport.pdf>).
- 116.) Maragou, N., Makri, A., Lampi, E., Thomaidis, N., Koupparis, M., "Migration of bisphenol A from polycarbonate baby bottles under real use conditions," *Food Additives and Contaminants*, Vol. 25, issue 3, 2008, pp. 373-383.
- 117.) Biedermann-Brem, S., Grob, K., Fjeldal, P., "Release of bisphenol A from polycarbonate baby bottles: mechanisms of formation and investigation of worst case scenarios," Norwegian Food Safety Authority, 2007, accessible at http://matportalen.no/Filer/fil_071101_Baby_bottels.pdf.
- 118.) Cited in the report as: Ehlert, K., Beumer, C., Groot, M., "Migration study of bisphenol A into water from polycarbonate baby bottles during microwave heating", 2008 (in press).
- 119.) Miyamoto, K., Kotake, M., "Estimation of daily bisphenol A intake of Japanese individuals with emphasis on uncertainty and variability", *Environmental Science*, Vol. 13, Issue 1, 2006, pp.15-29.
- 120.) Le, H., Carlson, E., Chue, J., Belcher, S., "Bisphenol A is released from polycarbonate drinking bottles and mimics the neurotoxic actions of estrogen in developing cerebellar neurons," *Toxicology Letters*, Vol. 176, 2008, pp. 149-156.
- 121.) CFSAN's "Infant formula" page at www.cfsan.fda.gov/~dms/inf-toc.html.

122.) Mead Johnson web site:

www.meadjohnson.com/app/iwp/HCP/Content2.do?dm=mj&id=/HCP_Home/Product_Information/Instructions_for_Preparation.

123.) Fein, S. B., Falci, C. D., 1999, "Infant Formula Preparation, Handling, and Related Practices in the United States", *Journal of the American Dietetic Association*, Vol, 99, No. 10, p. 1234.

124.) The Fein publication referenced Schuman, A, "Pediatrics 1990: Facts and fantasies, myths and misconceptions", *Clinical Pediatrics*, Vol. 29, 1990, pp. 558-564. (see Fein publication above)

125.) Available at www.who.int/foodsafety/publications/micro/pif2007/en/. "Pediatrics for Medical Students," 2nd Edition, D. Bernstein and S. P. Shelov, Lippincott, Williams & Wilkins, Published 2003 (ISBN 0781729416), Ch. 4, p. 88.

126.) "Caring for Your Baby & Young Child: from Birth to Age 5," S. Shelov, Oxford University Press, 1998 (ISBN 0192627783), Ch. 4, p. 101.

127.) Available at www.eatright.org/cps/rde/xchg/ada/hs.xsl/nutrition_1562_ENU_HTML.htm. "Preparing baby formula" by V. Iannelli, MD on About.com: Pediatrics (http://pediatrics.about.com/od/weeklyquestion/a/0707_bby_formla.htm).

128.) "Heading home with your newborn from Birth to Reality", Jana, L. and Shu, J., American Academy of Pediatrics, 2005.

129.) www.fda.gov/ohrms/dockets/AC/03/slides/3939s1_Anderson.ppt

130.) www.fda.gov/ohrms/dockets/ac/03/briefing/3939b1_tab4c_coversheet.pdf

131.) Cho, C.Y., Shin, B.S., Jung, J.H., Kim, D.H., Lee, K.C., Han, S.Y., Kim, H.S., Lee, B.M., Yoo, S.D. (2002) Pharmacokinetic scaling of bisphenol A by species-invariant time methods. *Xenobiotica* 32(10):925-934

132.) Csanády, G.A., Oberste-Frielinghaus, H.R., Semder, B., Baur, C., Schneider, K.T., Filser, B. (2002) Distribution and unspecific protein binding of the xenoestrogens bisphenol A and daidzein. *Arch. Toxicol.* 76:299-305

133.) Domoradzki, JY, Thornton, CM, Pottenger, LH, Hansen, SC, Card, TL, Markham, DA, Dryzga, MD, Shiotsuka, RN and Waechter, JM. (2004) Age and dose dependency of the pharmacokinetics and metabolism of bisphenol A in neonatal Sprague Dawley rats following oral administration . *Toxicol. Sci.* 77:230-242.

134.) Domoradzki, JY, Pottenger, LH, Thornton, CM, Hansen, SC, Card, TL, Markham, DA, Dryzga, MD, Shiotsuka, RN and Waechter, JM (2003). Metabolism and pharmacokinetics of bisphenol A (BPA) and the embryo fetal distribution of BPA and BPA monoglucuronide in CD Sprague Dawley rats at three gestational stages. *Toxicol. Sci.* 76:21-34.

135.) Elsby, R., Maggs, J.L., Ashby, J., Park, B.K. (2001) Comparison of the modulatory effects of human and rat liver microsomal metabolism on the estrogenicity of bisphenol A: implications for extrapolation to humans. *J. Pharmacol. Exp. Therap.* 297:103-113

- 136.) Hutter, J.C., Luu, H.M., Kim, C.S. (2004) A dynamic simulation of bisphenol A dosimetry in neuroendocrine organs. *Toxicol. Indust. Health* 20:29-40
- 137.) Iguchi T. (1992) Cellular effects of early exposure to sex hormones and antihormones. *Int. Rev. Cytol.* 139:1-57
- 138.) Inoue, H, Yuki, G, Yokota, H and Kato, S. (2003) Bisphenol A glucuronidation and absorption in rat intestine. *Drug Metab. Dispos.* 31:140 144.
- 139.) Inoue, H, Tsuruta, A, Kudo, S, Ishii T, Fukukushima, Y, Iwano, H, Yokota, H and Kato, S. (2004) Bisphenol A glucuronidation and excretion in liver of pregnant and nonpregnant female rats. *Drug Metab. Dispos.* 33:55 59.
- 140.) Kim, YH, Kim CS, Park, S, Han, SY, Pyo, MY and Yang M. (2003) Gender differences in the levels of bisphenol A metabolites in urine. *Biochem. Biophys. Res. Commun.* 312:441 448
- 141.) Kim, C.S., Sapienza, P.P., Ross, I.A., Johnson, W., Luu, H.M.D., Hutter, J.C. (2004) Distribution of bisphenol A in the neuroendocrine organs of female rats. *Toxicol. Indust. Health* 20:41-50
- 142.) Kurebayashi, H, Harada, R, Stewart, RK, Numata, H and Ohno, Y. (2002) Disposition of a low dose of bisphenol A in male and female cynomolgus monkeys. *Toxicol. Sci.* 69:32 42.
- 143.) Kurebayashi, H, Betsui, H and Ohno, Y. (2003) Disposition of a low dose of ¹⁴C bisphenol A in male rats and its main biliary excretion as BPA glucuronide. *Toxicol. Sci.* 73:17 25.
- 144.) Kurebayashi, H, Nagatsuka, S, Nemoto, H, Noguchi, H and Ohno, Y. (2005) Disposition of low doses of ¹⁴C bisphenol A in male, female, pregnant, fetal, and neonatal rats. *Arch. Toxicol.* 79:243 252.
- 145.) Kawade, N., Onishi, S. (1981) The prenatal and postnatal development of UDP-glucuronyl transferase activity towards bilirubin and the effect of premature birth on this activity in human liver. *Biochem. J.* 196:257-260
- 146.) Matsumoto, J, Yokota, H and Yuasa, A. (2002) Developmental increases in rat hepatic microsomal UDP glucuronosyl transferase activities toward xenoestrogens and decreases during pregnancy. *Environ. Health Perspect.* 110:193 196.
- 147.) Miyakoda, H, Tabata, M, Onodera, S and Takeda, K. (1999) Passage of bisphenol A into the fetus of the pregnant rat. *Journal of Health Science* 45:318 23.
- 148.) Miyakoda, H, Tabata, M, Onodera, S and Takeda, K. (2000) Comparison of conjugative activity, conversion of bisphenol A to bisphenol A glucuronide, in fetal and mature male rat. *Journal of Health Science* 46:269 274.
- 149.) Negishi, T, Tominaga, T, Ishii, Y, Kyuwa, S, Hayasaka, I, Kuroda, Y and Yoshikawa, Y (2004) Comparative study on toxicokinetics of bisphenol A in F344 rats, monkeys, (*Macaca fascicularis*), and chimpanzees (*Pan troglodytes*). *Exp. Anim.* 53:391 394.
- 150.) Pottenger, LH, Domoradzki, JY, Markham, DA, Hansen, SC, Cagen, SZ and Waechter, JM. (2000) The relative bioavailability and metabolism of bisphenol A in rats is dependent upon the route of administration. *Toxicol. Sci.* 54:3 18.

- 151.) Prichett, JJ, Kuester, RK and Sipes, IG (2002) Metabolism of bisphenol a in primary cultured hepatocytes from mice, rats, and humans. *Drug. Metab. Dispos.* 30:1180-1185.
- 152.) Sakamoto, H, Yokota, H, Kibe, R, Sayama, Y and Yuasa, A. (2002) Excretion of bisphenol A glucuronide into the small intestine and deconjugation in the cecum of the rat. *Biochim. Biophys. Acta.* 1573:171-176.
- 153.) Schonfelder G, Wittfoht, W, Hopp, H, Talsness, CE, Paul, M, and Chahoud, I. (2002) Parent bisphenol A accumulation in the human maternal fetal placental unit. *Environ. Health Perspect.* 110:A703-A707.
- 154.) Shin, BS, Yoo, SD, Cho, CY, Jung, JH, Lee, BM, Kim, JH, Lee, KC, Han, SY, Kim, HS, and Park KL. (2004) Maternal fetal disposition of bisphenol a in pregnant Sprague Dawley rats. *J. Toxicol. Environ. Health A* 65:395-406.
- 155.) Snyder, RW, Maness, SC, Gaido, KW, Welsch, F, Sumner, SC and Fennell, TR. (2000) Metabolism and disposition of bisphenol A in female rats. *Toxicol. Appl. Pharmacol.* 168:225-234.
- 156.) Takahashi, O and Oishi, S. (2000) Disposition of orally administered 2,2 bis(4 hydroxyphenyl) propane (Bisphenol A) in pregnant rats and the placental transfer to fetuses. *Environ. Health Perspect.* 108:931-935.
- 157.) Taylor, J.A., Welshons, W.V., vom Saal, F.S. (2008) No effect of route of exposure (oral; subcutaneous injection) on plasma bisphenol A throughout 24 hr after administration in neonatal female mice. *Reprod. Toxicol.* 25:169-176
- 158.) Teeguarden, JG, Waechter, JM, Clewell, HJ, Covington, TR and Barton, HA (2005) Evaluation of oral and intravenous route pharmacokinetics, plasma protein binding, and uterine tissue dose metrics of bisphenol A: A physiological based pharmacokinetic approach. *Toxicol. Sci.* 85:823-838.
- 159.) Tominaga, T., Negishi, T., Hirooka, H., Miyachi, A., Inoue, A., Hayasaka, I., Yoshikawa, Y. (2006) Toxicokinetics of bisphenol A in rats, monkeys and chimpanzees by the LC-MS/MS method. *Toxicology* 226:208-217.
- 160.) Uchida, K, Suzuki, A, Kobayashi, Y, Buchanan, DL, Sato, T, Watanabe, H, Katsu, Y, Suzuki, J, Asaoka, K, Mori, C (2002) Bisphenol A administration during pregnancy results in fetal exposure in mice and monkeys. *Journal of Health Science* 48:579-582.
- 161.) Upmeyer, A, Degen, GH, Diel, P, Michna, H, and Bolt, HM (2000) Toxicokinetics of bisphenol in female DA/Han rats after a single i.v. and oral administration. *Arch. Toxicol.* 74:431-436
- 162.) Völkel, W, Colnot, T, Csanady, GA, Filser, JG and Dekant, W. (2002) Metabolism and kinetics of bisphenol a in humans at low doses following oral administration. *Chem. Res. Toxicol.* 15:1281-1287.
- 163.) Waechter, J, Domoradzki, J., Thornton, C., Markham, D. (2007) Factors affecting the accuracy of bisphenol A and bisphenol A-monoglucuronide estimates in mammalian tissues and urine samples. *Toxicol. Mech. Method.* 17:13-24
- 164.) Ye, X., Bishop, A., Reidy, J.A., Needham, L.L., Calafat, A.M. (2007) Temporal stability of the

conjugated species of bisphenol A, parabens, and other environmental phenols in human urine. *J. Exposure Science and Environmental Epidemiology* 17:567-572

165.) Yokota, H, Iwano, H, Endo, M, Kobayashi, T, Inoue, H, Ikushiro, SI and Yuasa, A. (1999) Glucuronidation of the environmental estrogen bisphenol A by an isoform of UDP glucuronosyl transferase, UGT2B1, in the rat liver. *Biochem. Journal* 340:405-409.

166.) Yoo, SD, Shin, BS, Kwack, SJ, Lee, BM, Park KL, Han SY and Kim, HS. (2000) Pharmacokinetic disposition of bisphenol A in rats after intravenous administration. *J. Toxicol. Environ. Health.* 60:131-139.

167.) Yoo, SD, Shin, BS, Lee, BM, Lee, KC, Han SY, Kim, HS, Kwack, SJ and Park KL. (2001) Bioavailability and mammary excretion of bisphenol A in Sprague Dawley rats. *J. Toxicol. Environ. Health.* 64:417-426.

168.) Zalko, D, Soto, AM, Dolo, L, Dorio, C, Rathahao, E, Debrauwer, L, Faure, R. and Cravedi, JP (2003) Biotransformation of bisphenol A in a mammalian model: answers and new questions raised by low dose metabolic fate studies in pregnant CD1 mice. *Environ. Health Perspect.* 111:309-319.

169.) Jones, C. April 17, 2008. Food Navigator USA. New BPA evidence piles pressure on FDA to set exposure limits. <http://www.foodnavigator~usa.com/news/printNewsBis.asp?id=84718>

170.) Eyre, C. April 3, 2008. Food Navigator USA. Pregnant women told to avoid BPA packaging. New BPA evidence piles pressure on FDA to set exposure limits. <http://www.foodnavigator~usa.com/news/printNewsBis.asp?id=84718>

171.) Ahmed, E. Dec. 6, 2007. Food Navigator USA. Bisphenol A in infant formula at 'dangerous' levels, says group. <http://www.foodnavigator~usa.com/news/printNewsBis.asp?id=81877>

172.) Ahmed, E. Aug. 1, 2007. FoodNavigator USA. Unsafe levels of bisphenol A found in humans, scientist say. <http://www.foodnavigator~usa.com/news/printNewsBis.asp?id=7879>

173.) Seaborg, E. July, 2007. *Endocrine News*. Does breast cancer start in the womb. Alarming findings in bisphenol A Studies. P. 12-14.

174.) Reynolds, G. March 8, 2007. Food Navigator USA. Cancer chemical found at high levels in baby food. <http://www.foodnavigator~usa.com/news/printNewsBis.asp?id=74834>

175.) Smith, M. June 1, 2006. Neonatal estrogen exposure leads to prostate cancer in rats. *MedPage Today*. <http://www.medpagetoday.com/tbprint.cfm?tbid=3443>

176.) Keri, R.A. *et al.* (2007). *Reproductive Toxicology*. An evaluation of evidence for the carcinogenic activity of bisphenol A. 24(2): 240-252.

177.) Yoshida, M. *et al.* (2004). *J. Reproduction and Development*. Maternal Exposure to low doses of bisphenol A has no effects on development of female reproductive tract and uterine carcinogenesis in Donryu rats. 50(3): 349-360.

178.) Vandenberg, L. N. *et al.* (2007). *Endocrinology*. Exposure to environmentally relevant doses of the xenoestrogen bisphenol A alters development of the fetal mouse mammary gland. 148(1): 116-127.

- 179.) Munoz-de-Toro, M. *et al.* (2005). *Endocrinology*. Perinatal exposure to bisphenol A alters peripubertal mammary gland development in mice. 146(9): 4138-4147.
- 180.) Welshons, W.V. *et al.* (2006). *Endocrinology*. Large effects from small exposures. III. Mechanisms mediating effects of bisphenol A at levels of human exposure. 147(6): Supplement: S56-S69.
- 181.) Fenton, S. E. (2006). *Endocrinology*. Endocrine-disrupting compounds and mammary gland development: Early exposure and later life consequences. 147(6): Supplement S18-S24.
- 182.) Ho, Shuk-Mei *et al.* (2006). *Cancer Research*. Developmental exposure to estradiol and bisphenol A increases susceptibility to prostate carcinogenesis and epigenetically regulates phosphodiesterase type 4 variant 4. 66(11): 5624- 5632.
- 183.) Prins, G. S. *et al.* (2008). *Fertility and Sterility*. Developmental exposure to bisphenol A increases prostate cancer susceptibility in adult rats: epigenetic mode of action is implicated. 89(1): e41.
- 184.) Health and Environmental Sciences Mammalian and Environmental Toxicology Research Laboratory, Dow Chemical USA, Completed 06/06/1985 (WA 2000-20, Task 01).
- 185.) International Research and Development Corporation, 08/03/1976 (WA 2000-20, Task 02).
- 186.) Mammalian and Environmental Toxicology Research Laboratory Health and Environmental Sciences, Dow Chemical USA, 03/18/1988 (WA 2000-20, Task 03).
- 187.) Kubo K, Arai O, Omura M, Watanabe R, Ogata R, Aou S (2003) *Neurosci Res*. Low dose effects of bisphenol A on sexual differentiation of the brain and behavior in rats. 45(3): 345-356.
- 188.) Teratologic Evaluation of Bisphenol A administered to CD 8 rats on gestation days 6-15 (NTP 85-089)
- 189.) Teratologic Evaluation of Bisphenol A administered to CD-1 mice on gestation days 6-15 (NTP 85-088)
- 190.) Bisphenol A A reproduction and fertility assessment m CD-1 mice when administered in the feed (NTP 85-192)
- 191.) Bisphenol A: A reproduction and fertility assessment in CD-1 mice when administered via subcutaneous silastic implants (NTP 84-105)
- 192.) Rubin BS, Murray MK, Damassa DA, King JC, Soto AM (2001) *Environ Health Perspect*. Perinatal exposure to low doses of bisphenol A affects body weight, patterns of estrous cyclicity, and plasma LH levels. 109(7): 675-680.
- 193.) Thayer KA, Foster PM (2007) *Environ Health Perspect*. Workgroup Report: National Toxicology Program workshop on hormonally induced reproductive tumors - relevance of rodent bioassays. 115(9): 1351-1356.
- 194.) Tinwell H, Haseman J, Lefevre PA, Wallis N, Ashby J (2002) *Toxicol Sci*. Normal sexual development of two strains of rat exposed in utero to low doses of bisphenol A. 68(2): 339-348.

- 195.) Gupta C (2000) *Proc Soc Exp Biol Med*. Reproductive malformation of the male offspring following maternal exposure to estrogenic chemicals. 224(2): 61-68.
- 196.) Ichihara T, Yoshino H, Imai N, Tsutsumi T, Kawabe M, Tamano S, Inaguma S, Suzuki S, Shirai T (2003) *J Toxicol Sci*. Lack of carcinogenic risk in the prostate with transplacental and lactational exposure to bisphenol A in rats. 28(3): 165-171.
- 197.) Kwon S, Stedman DB, Elswick BA, Cattley RC, Welsch F (2000) *Toxicol Sci*. Pubertal development and reproductive functions of Crl:CD BR Sprague-Dawley rats exposed to bisphenol A during prenatal and postnatal development. 55(2): 399-406.
- 198.) Nagel SC, vom Saal FS, Thayer KA, Dhar MG, Boechler M, Welshons WV (1997) *Environmental Health Perspectives*. Relative Binding Affinity-Serum Modified Access (RBA-SMA) Assay Predicts the Relative In Vivo Bioactivity of the Xenoestrogens Bisphenol A and Octylphenol. 105(1): 70-76.
- 199.) Richter CA, Taylor JA, Ruhlen RL, Welshons WV, Vom Saal FS (2007) *Environ Health Perspect*. Estradiol and Bisphenol A stimulate androgen receptor and estrogen receptor gene expression in fetal mouse prostate mesenchyme cells. 115(6): 902-908.
- 200.) Wang S, Garcia AJ, Wu M, Lawson DA, Witte ON, Wu H (2006) *Proc Natl Acad Sci U S A*. Pten deletion leads to the expansion of a prostatic stem/progenitor cell subpopulation and tumor initiation. 103(5): 1480-1485.
- 201.) Evaluation of Reproductive Organ Development in CF-1 Mice Following Prenatal Exposure to Bisphenol A. MPI Research Study ID 328-046. Sponsored by The Society of the Plastics Industry, Inc. Completed 10/09/1998, Submitted to FDA in FMF 580.
- 202.) F.S.vomSaal, B.G.Timms, M.M.Montano, P.Palanza, K.A.Thayer, S.C.Nagel, M.D.Dhar, V.K.Ganjam, S.Parmigiani, W.V.Welshons. Prostate enlargement in mice due to fetal exposure to low doses of estradiol or diethylstilbestrol and opposite effects at high doses, *Proceedings of the National Academy of Sciences of the United States of America*, 94, (1997) 2056-2061.
- 203.) F.S.vom Saal, P.S.Cooke, D.L.Buchanan, P.Palanza, K.A.Thayer, S.C.Nagel, S.Parmigiani, W.V.Welshons. A physiologically based approach to the study of bisphenol A and other estrogenic chemicals on the size of reproductive organs, daily sperm production, and behavior, *Toxicology and Industrial Health*, 14, (1998) 239-260.
- 204.) Part 1: Literature Review of Neurobehavioral Effects of Bisphenol A, Exponent, 1010 14th Street, San Francisco, California, dated June 10, 2005.
- 205.) Della Seta, D., Minder, I., Dessi-Fulgheri, F., and Farabollini, F. (2005) Bisphenol-A exposure during pregnancy and lactation affects maternal behavior in rats. *Brain Res. Bull.* 65:255-260.
- 206.) Facciolo, R.M., Madeo, M., Alò, R., Canonaco, M., and Dessi-Fulgheri, F. (2005) Neurobiological effects of bisphenol A may be mediated by somatostatin subtype 3 receptors in some regions of the developing brain. *Toxicol. Sci.* 88:477-484.
- 207.) Fujimoto, T., Kubo, K., and Aou, S. (2006) Prenatal exposure to bisphenol A impairs sexual differentiation of exploratory behavior and increases depression-like behavior in rats. *Brain Res.* 1068:49-55.

- 208.) Funabashi, T., Kawaguchi, M., Furuta, M., Fukushima, A., and Kimura, F. (2004) Exposure to bisphenol A during gestation and lactation causes loss of sex difference in corticotrophin-releasing hormone-immunoreactive neurons in the bed nucleus of the stria terminalis of rats. *PNEC* 29:475-485.
- 209.) Honma, T., Miyagawa, M., Suda, M., Wang, R.-S., Kobayashi, K., and Sekiguchi, S. (2006) Effects of perinatal exposure to bisphenol A on brain neurotransmitters in female rat offspring. *Indus. Health* 44:510-524.
- 210.) Ishido, M., Morita, M., Oka, S., Masuo, Y. (2005) Alteration of gene expression of G protein-coupled receptors in endocrine disruptors-caused hyperactive rats. *Reg. Peptides* 126:145-153.
- 211.) Ishido, M., Yomemoto, J., Morita, M. (2007) Mesencephalic neurodegeneration in the orally administered bisphenol A-caused hyperactive rats. *Toxicol. Lett.* 173:66-72.
- 212.) Kawai, K., Murakami, S., Senba, E., Yamanaka, T., Fujiwara, Y., Arimura, C., Nozaki, T., Takii, M., and Kubo, C. (2007) Changes in estrogen receptors α and β expression in the brain of mice exposed prenatally to bisphenol A. *Reg. Toxicol. Pharmacol.* 47:166-170.
- 213.) MacLusky, N., Hajszan, T., Leranth, C. (2005) The environmental estrogen bisphenol A inhibits estradiol-induced hippocampal synaptogenesis. *Environ. Health Persp.* 113:675-679.
- 214.) Masuo, Y., Morita, M., Oka, S., and Ishido, M. (2004a) Motor hyperactivity caused by a deficit in dopaminergic neurons and the effects of endocrine disruptors: a study inspired by the physiological roles of PACAP in brain. *Reg. Peptides* 123:225-234.
- 215.) Masuo, Y., Ishido, M., Morita, M., and Oka, S. (2004b) Effects of neonatal treatment with 6-hydroxydopamine and endocrine disruptors on motor activity and gene expression in rats. *Neural Plasticity* 11:59-76.
- 216.) Mizuo, K., Narita, M., Miyagawa, K., Narita, M., Okuno, E., and Suzuki, T. (2004) Prenatal and neonatal exposure to bisphenol-A affects the morphine-induced rewarding effect and hyperlocomotion in mice. *Neurosci. Lett.* 356:95-98.
- 217.) Nakamura, K., Itoh, K., Yaoi, T., Fujiwara, Y., Sugimoto, T., and Fushiki, S. (2006) Murine neocortical histogenesis is perturbed by prenatal exposure to low doses of bisphenol A. *J. Neurosci. Res.* 84:1194-1205.
- 218.) Nakamura, K., Itoh, K., Sugimoto, T., and Fushiki, S. (2007) Prenatal exposure to bisphenol A affects adult murine neocortical structure. *Neurosci. Lett.* 420:100-105.
- 219.) Narita, M., Miyagawa, K., Mizuo, K., Yoshida, T., and Suzuki, T. (2006) Prenatal and neonatal exposure to low-dose of bisphenol-A enhance the morphine-induced hyperlocomotion and rewarding effect. *Neurosci. Lett.* 402:249-252.
- 220.) Narita, M., Miyagawa, K., Mizuo, K., Yoshida, T., and Suzuki, T. (2007) Changes in central dopaminergic systems and morphine reward by prenatal and neonatal exposure to bisphenol-A in mice: evidence for the importance of exposure period. *Addict. Biol.* 12:167-172.

- 221.) Nishizawa, H., Morita, M., Sugimoto, M., Imanishi, S., and Manabe, N. (2005a) Effects of in utero exposure to bisphenol A on mRNA expression of arylhydrocarbon and retinoid receptors in murine embryos. *J. Reprod. Dev.* 51:315-324.
- 222.) Nishizawa, H., Imanishi, S., and Manabe, N. (2005b) Effects of exposure in utero to bisphenol A on the expression of aryl hydrocarbon receptor, related factors, and xenobiotic metabolizing enzymes in murine embryos. *J. Reprod. Dev.* 51:593-605.
- 223.) Patisaul, H.B., Fortino, A.E., and Polston, E.K. (2006) Neonatal genistein or bisphenol-A exposure alters sexual differentiation of the AVPV. *Neurotoxicol. Teratol.* 28:111-118.
- 224.) Patisaul, H.B., Fortino, A.E., and Polston, E.K. (2007) Differential disruption of nuclear volume and neuronal phenotype in the preoptic area by neonatal exposure to genistein and bisphenol-A. *Neurotoxicol.* 28:1-12.
- 225.) Porrini, S., Belloni, V., Della Seta, D., Farabollini, F., Giannelli, G., and Dessi-Fulgheri, F. (2005) Early exposure to a low dose of bisphenol A affects socio-sexual behavior of juvenile female rats. *Brain Res. Bull.* 65:261-266.
- 226.) Razzoli, M., Valsecchi, P., and Palanza P. (2005) Chronic exposure to low doses bisphenol A interferes with pair-bonding and exploration in female Mongolian gerbils. *Brain Res. Bull.* 65:249-254.
- 227.) Rubin, B.S., Lenkowski, J.R., Schaeberle, C.M., Vandenberg, L.N., Ronsheim, P.M., and Soto, A.M. (2006) Evidence of altered brain sexual differentiation in mice exposed perinatally to low, environmentally relevant levels of bisphenol A. *Endocrinol.* 147:3681-3691.
- 228.) Shikimi, H., Sakamoto, H., Mezaki, Y., Ukena, K., and Tsutsui, K. (2004) Dendritic growth in response to environmental estrogens in the developing Purkinje cell in rats. *Neurosci. Lett.* 364:114-118.
- 229.) Tando, S., Itoh, K., Yaoi, T., Ikeda, J., Fujiwara, Y., and Fushiki, S. (2007) Effects of pre-and neonatal exposure to bisphenol A on murine brain development. *Brain Develop.* 29:352-356.
- 230.) Xu, X., Liu, Y., Sadamatsu, M., Tsutsumi, S., Akaike, M., Ushijima, H., and Kato, N. (2007) Perinatal bisphenol A affects the behavior and SRC-1 expression of male pups but does not influence on the thyroid hormone receptors and its responsive gene. *Neurosci. Res.* 58:149-155.
- 231.) Zoeller, R.T., Bansal, R., Parris, C. (2005) Bisphenol-A, an environmental contaminant that acts as a thyroid hormone receptor antagonist in vitro, increases serum thyroxine, and alters RC3/neurogranin expression in the developing rat brain. *Endocrinol.* 146:607-612.
- 232.) Akingbemi, B.T., Sottas, C.M., Koulova, A.I., Klinefelter, G.R., and Hardy, M.P., 2004. Inhibition of testicular steroidogenesis by the xenoestrogen bisphenol a is associated with reduced pituitary luteinizing hormone secretion and decreased steroidogenic enzyme gene expression in rat Leydig cells. *Endocrinology* 145, 592-603.
- 233.) Choi, I.S., Cho, J.H., Park, E.J., Park, J.W., Kim, S.H., Lee, M.G., Choi, B.J., and Jang, I.S., 2007. Multiple effects of bisphenol A, an endocrine disrupter, on GABA(A) receptors in acutely dissociated rat CA3 pyramidal neurons. *Neuroscience Research* 59, 8-17.

- 234.) Facciolo, R.M., Alo, R., Madeo, M., Canonaco, M., and ssi-Fulgheri, F., 2002. Early cerebral activities of the environmental estrogen bisphenol A appear to act via the somatostatin receptor subtype sst(2). *Environmental Health Perspectives* 110, 397-402.
- 235.) Fini, J.B., Le Mevel, S., Turque, N., Palmier, K., Zalko, D., Cravedi, J.P., and Demeneix, B.A., 2007. An in vivo multiwell-based fluorescent screen for monitoring vertebrate thyroid hormone disruption. *Environmental Science & Technology* 41, 5908-5914.
- 236.) Funabashi, T., Kawaguchi, M., and Kimura, F., 2001. The endocrine disrupters butyl benzyl phthalate and bisphenol A increase the expression of progesterone receptor messenger ribonucleic acid in the preoptic area of adult ovariectomized rats. *Neuroendocrinology* 74, 77-81.
- 237.) Funabashi, T., Nakamura, T.J., and Kimura, F., 2004. p-nonylphenol, 4-tert-octylphenol and bisphenol A increase the expression of progesterone receptor mRNA in the frontal cortex of adult ovariectomized rats. *Journal of Neuroendocrinology* 16, 99-104.
- 238.) Ghisari, M. and Bonefeld-Jorgensen, E.C., 2005. Impact of environmental chemicals on the thyroid hormone function in pituitary rat GH3 cells. *Molecular and Cellular Endocrinology* 244, 31-41.
- 239.) Jung, K.K., Kim, S.Y., Kim, T.G., Kang, J.H., Kang, S.Y., Cho, J.Y., and Kim, S.H., 2007. Differential regulation of thyroid hormone receptor-mediated function by endocrine disruptors. *Archives of Pharmacal Research* 30, 616-623.
- 240.) Kiguchi M, Fujita S, Lee J, Shimizu N, Koshikawa N. 2007 Behavioral responses to methylphenidate and in rats exposed neonatally to bisphenol-A. *J Oral Sci.* 49(4): 311-318.
- 241.) Kim, K., Son, T.G., Kim, S.J., Kim, H.S., Kim, T.S., Han, S.Y., and Lee, J., 2007. Suppressive effects of bisphenol a on the proliferation of neural progenitor cells. *Journal of Toxicology and Environmental Health-Part A-Current Issues* 70, 1288-1295.
- 242.) Lee, Y.M., Seong, M.J., Lee, J.W., Lee, Y.K., Kim, T.M., Nam, S.Y., Kim, D.J., Yun, Y.W., Kim, T.S., Han, S.Y., and Hong, J.T., 2007. Estrogen receptor independent neurotoxic mechanism of bisphenol A, an environmental estrogen. *Journal of Veterinary Science* 8, 27-38.
- 243.) Leranth, C., Szigeti-Buck, K., MacLusky, N.J., and Hajszan, T., 2008. Bisphenol a prevents the synaptogenic response to testosterone in the brain of adult male rats. *Endocrinology* 149, 988-994.
- 244.) Nishizawa, H., Manabe, N., Morita, M., Sugimoto, M., Imanishi, S., and Miyamoto, H., 2003. Effects of In utero exposure to bisphenol A on expression of RAR alpha and RXR alpha mRNAs in murine embryos. *Journal of Reproduction and Development* 49, 539-545.
- 245.) Ogiue-Ikeda, M., Tanabe, N., Mukai, H., Hojo, Y., Murakami, G., Tsurugizawa, T., Takata, N., Kimoto, T., and Kawato, S., 2008. Rapid modulation of synaptic plasticity by estrogens as well as endocrine disrupters in hippocampal neurons. *Brain Research Reviews* 57, 363-375.
- 246.) Patisaul, H.B. and Bateman, H.L., 2008. Neonatal exposure to endocrine active compounds or an ER agonist increases adult anxiety and aggression in gonadally intact male rats. *Hormones and Behavior* 53, 580-588.

247.) Seiwa, C., Nakahara, J., Komiyama, T., Katsu, Y., Iguchi, T., and Asou, H., 2004. Bisphenol A exerts thyroid-hormone-like effects on mouse oligodendrocyte precursor cells. *Neuroendocrinology* 80, 21-30.

248.) Takagi, H., Shibutani, M., Masutomi, N., Uneyama, C., Takahashi, N., Mitsumori, K., and Hirose, M., 2004. Lack of maternal dietary exposure effects of bisphenol A and nonylphenol during the critical period for brain sexual differentiation on the reproductive/endocrine systems in later life. *Archives of Toxicology* 78, 97-105.

249.) FDA memorandum - Memorandum to the File (FAP 8T4773). Corrections to Bisphenol A Reviews. Twaroski/Lin, 08/12/2008.