

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ASPARLAS safely and effectively. See full prescribing information for ASPARLAS.

**ASPARLAS™ (calaspargase pegol - mknl) injection, for intravenous use**  
Initial U.S. Approval: 2018

### INDICATIONS AND USAGE

ASPARLAS is an asparagine specific enzyme indicated as a component of a multi-agent chemotherapeutic regimen for the treatment of acute lymphoblastic leukemia in pediatric and young adult patients age 1 month to 21 years. (1.1)

### DOSAGE AND ADMINISTRATION

- Recommended Dosage: 2,500 units/m<sup>2</sup> intravenously no more frequently than every 21 days. (2.1)
- See Full Prescribing Information for important details regarding dosing modifications and preparation and administration. (2.2, 2.3)

### DOSAGE FORMS AND STRENGTHS

Injection: 3,750 units/5 mL (750 units/mL) in a single-dose vial. (3)

### CONTRAINDICATIONS

- History of serious hypersensitivity reactions to pegylated L-asparaginase. (4)
- History of serious thrombosis during L-asparaginase therapy. (4)
- History of serious pancreatitis related to previous L-asparaginase treatment. (4)

- History of serious hemorrhagic events during previous L-asparaginase therapy. (4)
- Severe hepatic impairment. (4)

### WARNINGS AND PRECAUTIONS

- Hypersensitivity: Observe patients for one hour after administration. Discontinue ASPARLAS in patients with serious hypersensitivity reactions. (5.1)
- Pancreatitis: Discontinue ASPARLAS in patients with pancreatitis. Monitor blood glucose (5.2)
- Thrombosis: Discontinue ASPARLAS for severe or life-threatening thrombosis. Provide anticoagulation therapy as indicated.
- Hemorrhage: Discontinue ASPARLAS for severe or life-threatening hemorrhage. Evaluate for etiology and treat. (5.3, 5.4)
- Hepatotoxicity: Monitor for toxicity through recovery from cycle. (5.4)

### ADVERSE REACTIONS

The most common (incidence ≥10%) grade ≥ 3 adverse reactions were elevated transaminase, bilirubin increased, pancreatitis and abnormal clotting studies. (6.1)

### USE IN SPECIFIC POPULATIONS

Lactation: Advise women not to breastfeed. (8.2)

To report SUSPECTED ADVERSE REACTIONS, contact Servier Pharmaceuticals LLC at 857-895-4853 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 12/2018

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

- 1.1 Acute Lymphoblastic Leukemia (ALL)

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage
- 2.2 Dose Modifications
- 2.3 Preparation and Administration

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity
- 5.2 Pancreatitis
- 5.3 Thrombosis
- 5.4 Hemorrhage
- 5.5 Hepatotoxicity

### 6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Immunogenicity

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy

8.2 Lactation

8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

### 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### 14 CLINICAL STUDIES

- 14.1 Acute Lymphoblastic Leukemia (ALL)

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

---

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

#### 1.1 Acute Lymphoblastic Leukemia

ASPARLAS is indicated as a component of a multi-agent chemotherapeutic regimen for the treatment of acute lymphoblastic leukemia in pediatric and young adult patients age 1 month to 21 years.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Recommended Dosage

The recommended dose of ASPARLAS is 2,500 units/m<sup>2</sup> given intravenously no more frequently than every 21 days.

#### 2.2 Dose Modifications

Monitor patients at least weekly, with bilirubin, transaminases, glucose and clinical examinations until recovery from the cycle of therapy. If an adverse reaction should occur, modify treatment according to Table 1.

**Table 1. Dose Modifications**

Adverse Reaction	Severity*	Action
Infusion Reaction or Hypersensitivity Reaction	Grade 1	<ul style="list-style-type: none"><li>Reduce the infusion rate by 50%</li></ul>
	Grade 2	<ul style="list-style-type: none"><li>Interrupt the infusion of ASPARLAS</li><li>Treat the symptoms</li><li>When symptoms resolve, resume the infusion and reduce the infusion rate by 50%</li></ul>
	Grade 3 to 4	<ul style="list-style-type: none"><li>Discontinue ASPARLAS permanently</li></ul>
Hemorrhage	Grade 3 to 4	<ul style="list-style-type: none"><li>Hold ASPARLAS.</li><li>Evaluate for coagulopathy and consider clotting factor replacement as needed.</li><li>Resume ASPARLAS with the next scheduled dose if bleeding is controlled.</li></ul>
Pancreatitis	Grades 3 to 4	<ul style="list-style-type: none"><li>Hold ASPARLAS for elevations in lipase or amylase &gt;3 times the ULN until enzyme levels stabilize or are declining</li><li>Discontinue ASPARLAS permanently if clinical pancreatitis is confirmed.</li></ul>
Thromboembolism	Uncomplicated deep vein thrombosis	<ul style="list-style-type: none"><li>Hold ASPARLAS.</li><li>Treat with appropriate antithrombotic therapy</li><li>Upon resolution of symptoms consider resuming ASPARLAS, while continuing antithrombotic therapy.</li></ul>
	Severe or life-threatening thrombosis	<ul style="list-style-type: none"><li>Discontinue ASPARLAS permanently.</li><li>Treat with appropriate antithrombotic therapy.</li></ul>
Hepatotoxicity	Total bilirubin more than 3 times to no more than 10 times the upper limit of normal	<ul style="list-style-type: none"><li>Hold ASPARLAS until Total bilirubin levels go down to ≤ 1.5 times the upper limit of normal</li></ul>

	Total bilirubin more than 10 times the upper limit of normal	<ul style="list-style-type: none"> <li>Discontinue ASPARLAS and do not make up for missed doses</li> </ul>
--	--	--

\*Grade 1 is mild, grade 2 is moderate, grade 3 is severe, and grade 4 is life-threatening

### 2.3 Preparation and Administration

ASPARLAS is a clear and colorless solution. Visually inspect parenteral drug products for particulate matter, cloudiness, or discoloration prior to administration. If any of these are present, discard the vial. Do not administer if ASPARLAS has been shaken or vigorously agitated, frozen, or stored at room temperature for more than 48 hours.

- Dilute ASPARLAS in 100 mL of 0.9% Sodium Chloride Injection, USP or 5% Dextrose Injection, USP using sterile/aseptic technique. Discard any unused portion left in a vial.
- After dilution, administer immediately into a running infusion of either 0.9% sodium chloride or 5% dextrose, respectively.
- Administer the dose over a period of 1 hour.
- Do not infuse other drugs through the same intravenous line during administration of ASPARLAS.
- The diluted solution may be stored for up to 4 hours at room temperature (15°C to 25°C [59°F to 77°F]) or refrigerated at 2°C to 8°C (36°F to 46°F) for up to 24 hours.
- Protect from light. Do not shake or freeze.

## 3 DOSAGE FORMS AND STRENGTHS

Injection: 3,750 units/5 mL (750 units/mL) clear, colorless solution in a single-dose vial.

## 4 CONTRAINDICATIONS

ASPARLAS is contraindicated in patients with:

- History of serious hypersensitivity reactions, including anaphylaxis, to pegylated L-asparaginase therapy [see [Warnings and Precautions \(5.1\)](#)];
- History of serious thrombosis during previous L-asparaginase therapy [see [Warnings and Precautions \(5.3\)](#)];
- History of serious pancreatitis during previous L-asparaginase therapy [see [Warnings and Precautions \(5.2\)](#)];
- History of serious hemorrhagic events during previous L-asparaginase therapy [see [Warnings and Precautions \(5.4\)](#)];
- Severe hepatic impairment [see [Warnings and Precautions \(5.5\)](#)].

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Hypersensitivity

Grade 3 and 4 hypersensitivity reactions including anaphylaxis have been reported in clinical trials with ASPARLAS with an incidence between 7 to 21% [see [Contraindications \(4\)](#), [Adverse Reactions \(6.1\)](#)]. Hypersensitivity reactions observed with other asparaginases include angioedema, lip swelling, eye swelling, erythema, blood pressure decreased, bronchospasm, dyspnea, pruritus and rash.

Because of the risk of serious allergic reactions (e.g., life-threatening anaphylaxis), administer ASPARLAS in a clinical setting with resuscitation equipment and other agents necessary to treat anaphylaxis (e.g., epinephrine, oxygen, intravenous steroids, antihistamines) [see [Dosage and Administration \(2\)](#)] and observe patients for 1 hour after administration. Discontinue ASPARLAS in patients with serious hypersensitivity reactions.

### 5.2 Pancreatitis

Cases of pancreatitis have been reported in clinical trials with ASPARLAS with an incidence between 12 to 16% [see [Adverse Reactions \(6.1\)](#)]. Hemorrhagic or necrotizing pancreatitis have been reported with other asparaginases.

Inform patients of the signs and symptoms of pancreatitis, which, if left untreated, could be fatal. Assess serum amylase and/or lipase levels to confirm early signs of pancreatic inflammation. Discontinue ASPARLAS if pancreatitis is suspected; if pancreatitis is confirmed, do not resume ASPARLAS [see [Dosage and Administration \(2.2\)](#)].

### 5.3 Thrombosis

Serious thrombotic events, including sagittal sinus thrombosis, have been reported in clinical trials with ASPARLAS with an incidence of 9 to 12%. Discontinue ASPARLAS in patients experiencing serious thrombotic events [see [Dosage and Administration \(2.2\)](#), [Adverse Reactions \(6.1\)](#)].

### 5.4 Hemorrhage

Hemorrhage associated with increased prothrombin time (PT), increased partial thromboplastin time (PTT), and hypofibrinogenemia have been reported in patients receiving ASPARLAS [see [Adverse Reactions \(6.1\)](#)]. Evaluate patients with signs and symptoms of hemorrhage with coagulation parameters including PT, PTT, fibrinogen. Consider appropriate replacement therapy in patients with severe or symptomatic coagulopathy [see [Dosage and Administration \(2.2\)](#)].

### 5.5 Hepatotoxicity

Hepatotoxicity and abnormal liver function, including elevations of transaminase, bilirubin (direct and indirect), reduced serum albumin, and plasma can occur. Evaluate bilirubin and transaminases at least weekly, during cycles of treatment that include ASPARLAS through at least 6 weeks after the last dose of ASPARLAS. In the event of serious liver toxicity, discontinue treatment with ASPARLAS and provide supportive care [see [Dosage and Administration \(2.2\)](#), [Contraindications \(4\)](#), [Adverse Reactions \(6.1\)](#)].

## 6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described in greater detail in other sections of the labeling:

- Hypersensitivity [see [Warnings and Precautions \(5.1\)](#)].
- Pancreatic Toxicity [see [Warnings and Precautions \(5.2\)](#)].
- Thrombosis [see [Warnings and Precautions \(5.3\)](#)].
- Hemorrhage [see [Warnings and Precautions \(5.4\)](#)].
- Hepatotoxicity [see [Warnings and Precautions \(5.5\)](#)].

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of ASPARLAS was investigated in Study DFCI 11-001, an open-label, randomized, active-controlled multicenter clinical trial that treated 237 children and adolescents with newly-diagnosed ALL or lymphoblastic lymphoma, with ASPARLAS 2,500 U/m<sup>2</sup> (n=118) or pegaspargase 2,500 U/m<sup>2</sup> (n=119) as part of a Dana Farber Cancer Institute (DFCI) ALL Consortium backbone therapy. The median age on enrollment was 5 years (range, 1-20) years. The majority of patients were male (62%) and white (70%). Most patients were considered standard risk (SR, 59%) and had B-cell lineage ALL (87%).

The median number of doses during the study was 11 doses for ASPARLAS (administered every three weeks) and 16 doses for pegaspargase (administered every two weeks). The median duration of exposure was 8 months for both ASPARLAS and pegaspargase.

There was 1 fatal adverse reaction (multi-organ failure in the setting of chronic pancreatitis associated with a pancreatic pseudocyst).

**Table 2** summarizes the incidence of selected Grades  $\geq 3$  adverse reactions that occurred in 2 or more patients receiving ASPARLAS. Because not all grade 1 and 2 adverse reactions were collected prospectively, only grade 3 and 4 adverse reactions are presented in Table 1.

**Table 2: Selected Grades ≥ 3 Adverse Reactions in Patients Receiving ASPARLAS With Multi-Agent Chemotherapy (Study DFCI 11-001)\***

Adverse Reaction†	ASPARLAS 2,500 U/m <sup>2</sup> N=118	Pegaspargase 2,500 U/m <sup>2</sup> N=119
	Grades ≥ 3 n (%)§	Grades ≥ 3 n (%)§
Elevated transaminase	61 (52)	79 (66)
Bilirubin increased	24 (20)	30 (25)
Pancreatitis	21 (18)	29 (24)
Abnormal clotting studies	17 (14)	25 (21)
Diarrhea	10 (9)	6 (5)
Hypersensitivity	9 (8)	8 (7)
Embolic and thrombotic events	9 (8)	10 (8)
Sepsis	6 (5)	7 (6)
Dyspnea	5 (4)	1 (1)
Hemorrhages	5 (4)	5 (4)
Fungal infection	4 (3)	3 (3)
Pneumonia	4 (3)	8 (7)
Arrhythmia	2 (2)	1 (1)
Cardiac failure	2 (2)	1 (1)

\* ASPARLAS and pegaspargase were administered as a component of multi-agent chemotherapy regimens.

† Grouped terms: **Elevated transaminase**: Alanine aminotransferase increased, Aspartate aminotransferase increased, Transaminases increased; **Bilirubin increased**: Bilirubin conjugated increased, Blood bilirubin increased; **Pancreatitis**: Amylase increased, Lipase increased, Pancreatic necrosis, Pancreatitis, Pancreatitis relapsing; **Abnormal clotting studies**: Activated partial thromboplastin time prolonged, Blood fibrinogen decreased; **Diarrhea**: Colitis, Diarrhea, Enterocolitis, Neutropenic colitis; **Hypersensitivity**: Anaphylactic reaction, Drug hypersensitivity, Hypersensitivity; **Embolic and thrombotic events SMQ**: Device related thrombosis, Disseminated intravascular coagulation, Embolism, Intracardiac thrombus, Intracranial venous sinus thrombosis, Pulmonary embolism, Superior sagittal sinus thrombosis, Thrombosis in device, Venous thrombosis, Venous thrombosis limb; **Sepsis**: Bacterial sepsis, Sepsis; **Dyspnea**: Hypoxia, Respiratory failure; **Hemorrhages SMQ** (excludes laboratory terms): Disseminated intravascular coagulation, Epistaxis, Hematoma, Hemorrhage intracranial, Melena, Esophageal ulcer hemorrhage, Small intestinal hemorrhage, Upper gastrointestinal hemorrhage; **Fungal infection**: Fungal infection, Hepatic infection fungal, Respiratory tract infection fungal, Splenic infection fungal, Systemic candida; **Pneumonia**: Lung infection, Pneumonia, Pneumonitis; **Arrhythmia**: Atrioventricular block complete, Sinus tachycardia, Ventricular arrhythmia; **Cardiac failure**: Ejection fraction decreased, Left ventricular dysfunction.

§ Grading is based on the Common Terminology Criteria for Adverse Events (CTCAE) v4.0.

In the subgroup of patients with B-cell lineage ALL, the complete remission rate in the ASPARLAS arm was 98% (95/97), compared to 99% in the pegaspargase arm; the Kaplan-Meier estimates of overall survival of the treatment arms were comparable.

#### Study AALL07P4

The safety of ASPARLAS was also evaluated in Study AALL07P4, an open-label, randomized, active-controlled, multicenter clinical trial that treated patients with newly-diagnosed high-risk B-precursor ALL using ASPARLAS 2,500 U/m<sup>2</sup> (n=43) or 2,100 U/m<sup>2</sup> (n=68), or pegaspargase 2,500 U/m<sup>2</sup> (n=52), as a

component of an augmented Berlin-Frankfurt-Münster (BFM) therapy regimen. The median age was 11 years (range 1 to 26 years); the median duration of exposure was 7 months for both ASPARLAS and pegaspargase. In this study, the induction mortality of patients treated with ASPARLAS was 2.8% (3 out of 111); there were no induction deaths among 52 patients treated with pegaspargase.

## **6.2 Immunogenicity**

As with all therapeutic proteins, there is potential for immunogenicity.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no available data on the use of ASPARLAS in pregnant women to inform a drug-associated risk of major birth defects and miscarriage. Published literature studies in pregnant animals suggest asparagine depletion may cause harm to the animal offspring (see [Data](#)). Advise patients of the potential risk to a fetus.

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies are 2%-4% and 15%-20%, respectively.

#### Data

##### *Animal Data*

Animal reproduction studies have not been conducted with ASPARLAS to evaluate its effect on reproduction and fetal development. Published literature studies in which pregnant rabbits were administered L-asparaginase or pregnant rats were deprived of dietary asparagine suggested harm to the animal offspring.

### **8.2 Lactation**

#### Risk Summary

There are no data on the presence of calaspargase pegol-mknl in human milk, the effects on the breastfed child, or the effects on milk production. Because many drugs are excreted in human milk and because of the potential for adverse reactions in a breastfed child, advise lactating women not to breastfeed while receiving ASPARLAS and for 3 months after the last dose.

### **8.3 Females and Males of Reproductive Potential**

Based on published literature studies in pregnant animals, ASPARLAS can cause fetal harm when administered to a pregnant woman [see [Use in Specific Populations \(8.1\)](#)].

#### Pregnancy Testing

Conduct pregnancy testing in females of reproductive potential prior to starting treatment with ASPARLAS.

#### Contraception

Advise females of reproductive potential to avoid becoming pregnant while receiving ASPARLAS. Females should use effective contraceptive methods, including a barrier method, during treatment and for at least 3 months after the last dose of ASPARLAS. Since there is a potential for an indirect interaction between ASPARLAS and oral contraceptives, the concomitant use of ASPARLAS and oral contraceptives is not recommended. Another, non-oral contraceptive method should be used in women of childbearing potential.

### **8.4 Pediatric Use**

The safety and effectiveness of ASPARLAS in the treatment of ALL have been established in pediatric patients 1 month to < 17 years (no data for the age group < 1 month old). Use of ASPARLAS in these age groups is supported by evidence from an adequate and well-controlled trial with additional safety

from a second trial. The trials included 208 children with ALL or lymphoblastic lymphoma treated with ASPARLAS; there were 19 infants (1 month to < 2 years old), 128 children (2 years to < 12 years old), and 61 adolescents (12 years to < 17 years old). There were no clinically meaningful differences in safety or nadir serum asparaginase activity across age groups [see [Adverse reactions \(6.1\)](#), [Clinical Studies \(14\)](#)].

## 11 DESCRIPTION

Calaspargase pegol-mknl contains an asparagine specific enzyme derived from *Escherichia coli*, as a conjugate of L-asparaginase (L-asparagine amidohydrolase) and monomethoxypolyethylene glycol (mPEG) with a succinimidyl carbonate (SC) linker. The SC linker is a chemically stable carbamate bond between the mPEG moiety and the lysine groups of L-asparaginase.

L-asparaginase is a tetrameric enzyme that is produced endogenously by *E. coli* and consists of identical 34.5 kDa subunits. Approximately 31 to 39 molecules of SC-PEG are linked to L-asparaginase; the molecular weight of each SC-PEG molecule is about 5 kDa. The activity of ASPARLAS is expressed in units (U).

ASPARLAS injection is supplied as a clear, colorless, preservative-free, isotonic sterile solution in phosphate-buffered saline, pH 7.3 that requires dilution prior to intravenous infusion. Each vial of ASPARLAS contains 3,750 units in 5 mL of solution. Each milliliter contains 750 units of calaspargase pegol-mknl; dibasic sodium phosphate, USP (5.58 mg); monobasic sodium phosphate, USP (1.20 mg); and sodium chloride, USP (8.50 mg) in water for injection, USP.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

L-asparaginase is an enzyme that catalyzes the conversion of the amino acid L-asparagine into aspartic acid and ammonia. The pharmacological effect of ASPARLAS is thought to be based on selective killing of leukemic cells due to depletion of plasma L-asparagine. Leukemic cells with low expression of asparagine synthetase have a reduced ability to synthesize L-asparagine, and therefore depend on an exogenous source of L-asparagine for survival.

### 12.2 Pharmacodynamics

Calaspargase pegol-mknl pharmacodynamic (PD) response was assessed through measurement of plasma and cerebrospinal fluid (CSF) asparagine concentrations *via* an LC-MS/MS assay.

Asparagine concentrations in plasma (N=41) were maintained below the assay limit of quantification for more than 18 days following a single dose of ASPARLAS 2,500 U/m<sup>2</sup> during the induction phase. Mean CSF asparagine concentrations decreased from a pretreatment concentration of 0.8 µg/mL (N=10) to 0.2 µg/mL on Day 4 (N=37) and remained decreased at 0.2 µg/mL (N=35) 25 days after the administration of a single dose of ASPARLAS 2,500 U/m<sup>2</sup> in the induction phase.

### 12.3 Pharmacokinetics

Calaspargase pegol-mknl pharmacokinetics (PK) were assessed through measurement of plasma asparaginase activity *via* a coupled enzymatic assay.

The plasma asparaginase activity pharmacokinetics were characterized in 43 patients (1 to 26 years) with newly diagnosed high risk B-precursor ALL treated with a multidrug backbone therapy. **Table 3** summarizes the plasma asparaginase activity pharmacokinetic parameters after a single dose of ASPARLAS 2,500 U/m<sup>2</sup> in the induction phase.

**Table 3: Plasma Asparaginase Activity Pharmacokinetic Parameters After a Single Dose of ASPARLAS 2,500 U/m<sup>2</sup> in Patients with ALL in Study AALL07P4**

Parameter	Arithmetic Mean (%CV) N=43
<b>General</b>	
C <sub>max</sub> (U/mL)	1.62 (23.0)
AUC <sub>0-25day</sub> (day·U/mL)	16.9 (23.2)*
AUC <sub>0-∞</sub> (day·U/mL) <sup>†</sup>	25.5 (30.4)*
<b>Absorption</b>	
T <sub>max</sub> (h) <sup>†</sup>	1.17 (1.05, 5.47) <sup>‡</sup>
<b>Distribution</b>	

V <sub>ss</sub> (L)	2.96 (84.3)*
<b>Elimination</b>	
t <sub>1/2</sub> (day) <sup>§</sup>	16.1 (51.9)*
Clearance (L/day)	0.147 (76.1)*

\* N= 42 evaluable subjects.

† T<sub>max</sub> generally near end of a 1 hour calaspargase pegol-mknl intravenous (IV) infusion.

‡ Median (10<sup>th</sup>, 90<sup>th</sup> percentiles).

§ Plasma asparaginase activity pharmacokinetics are nonlinear following ASPARLAS administration.

### Specific Populations

The impact of renal and hepatic impairment on the PK of calaspargase pegol-mknl is unknown.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Carcinogenicity, mutagenicity, and impairment of fertility studies have not been conducted with calaspargase pegol-mknl.

## **14 CLINICAL STUDIES**

### **14.1 Acute Lymphoblastic Leukemia (ALL)**

The determination of efficacy was based on a demonstration of the achievement and maintenance of nadir serum asparaginase activity (NSAA) above the level of 0.1 U/mL using ASPARLAS 2500 U/m<sup>2</sup> intravenously every 3 weeks. The pharmacokinetics of ASPARLAS were studied when used in combination with multiagent chemotherapy in 124 patients with B cell lineage acute lymphoblastic leukemia (ALL). Among these patients, the median age was 11.5 years (range 1 – 26); 62 (50%) were male, 102 (82%) white, 6 (5%) Asian, 5 (4%) Black or African American, 2 (2%) Native Hawaiian or Pacific Islander and 9 (7%) other or unknown. The results showed that 123 (99%, 95% CI: 96% - 100%) of the 124 patients maintained NSAA > 0.1 U/mL at weeks 6, 12, 18, 24 and 30.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

ASPARLAS injection is supplied as a clear, colorless, preservative-free sterile solution in single-dose vials containing 3,750 units of calaspargase pegol-mknl per 5 mL solution (NDC 72694-515-01).

Store ASPARLAS refrigerated at 2°C to 8°C (36°F to 46°F) in the original carton to protect from light. Do not shake or freeze product. Unopened vials may be stored at room temperature (15°C to 25°C [59°F to 77°F]) for no more than 48 hours.

## **17 PATIENT COUNSELING INFORMATION**

Advise Patients/Caregivers of the following risks of ASPARLAS:

### Hypersensitivity

Inform patients on the possibility of serious allergic reactions, including anaphylaxis. Instruct the patient on the symptoms of allergic reactions and to seek medical advice immediately if they experience such symptoms [see [Warnings and Precautions \(5.1\)](#)].

### Pancreatitis

Instruct patients on the signs and symptoms of pancreatitis and to seek immediate medical attention if they experience severe abdominal pain [see [Warnings and Precautions \(5.2\)](#)].

Instruct patients on the risk of hyperglycemia and glucose intolerance. Advise patients to seek medical advice if they experience excessive thirst or any increase in the volume or frequency of urination [see [Dosage and Administration \(2.2\)](#)].

### Thrombosis

Instruct patients on the risk of thrombosis and to seek medical advice immediately if they experience severe headache, arm or leg swelling, shortness of breath, or chest pain [see [Warnings and Precautions \(5.3\)](#)].



### Hemorrhage

Advise patients to report any unusual bleeding or bruising to their physician [see [Warnings and Precautions \(5.4\)](#)].

### Hepatotoxicity

Advise patients to contact their healthcare provider immediately for jaundice, severe nausea or vomiting, or easy bruising or bleeding [see [Warnings and Precautions \(5.5\)](#)].

### Pregnancy and Lactation

Advise female patients of reproductive potential to use effective contraceptive methods while receiving ASPARLAS and for at least 3 months after the last dose. Advise patients to notify their healthcare provider immediately in the event of a pregnancy or if pregnancy is suspected during ASPARLAS treatment. Since there is a potential for an indirect interaction between ASPARLAS and oral contraceptives, the concomitant use of ASPARLAS and oral contraceptives is not recommended [see [Use in Specific Populations \(8.3\)](#)].

Advise lactating women not to breastfeed during treatment with ASPARLAS and for at least 3 months after the last dose [see [Use in Specific Populations \(8.2\)](#)].

---

Manufactured by:

Servier Pharmaceuticals LLC.  
Boston, MA 02116  
U.S. License No. 2125

Copyright © 2018  
All rights reserved.

ASPARLAS™ is a registered trademark of Servier IP UK Ltd, a wholly owned indirect subsidiary of Les Laboratoires Servier. Servier Pharmaceuticals LLC is a wholly own subsidiary of Les Laboratoires Servier.