

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ASACOL HD safely and effectively. See full prescribing information for ASACOL HD.

ASACOL[®] HD (mesalamine) delayed-release tablets, for oral use
Initial U.S. Approval: 1987

INDICATIONS AND USAGE

Asacol HD is an aminosalicilate indicated for the treatment of moderately active ulcerative colitis in adults. (1)

Limitation of Use:

Safety and effectiveness of Asacol HD beyond 6 weeks have not been established (1)

DOSAGE AND ADMINISTRATION

Important Administration Instructions (2.1):

- Do not substitute one Asacol HD 800 tablet for two mesalamine delayed-release 400 mg oral products (2.1)
- Evaluate renal function prior to initiation of Asacol HD (2.1, 5.1)
- Take on an empty stomach, at least 1 hour before and 2 hours after a meal
- Swallow whole; do not cut, break or chew the tablets.

Treatment of Moderately Active Ulcerative Colitis (2.2):

- Recommended dosage is 1600 mg (two 800 mg tablets) three times daily for 6 weeks.

DOSAGE FORMS AND STRENGTHS

Delayed-release tablets: 800 mg (3)

CONTRAINDICATIONS

Known or suspected hypersensitivity to salicylates or aminosalicylates or to any of the ingredients of Asacol HD tablets (4, 5.3)

WARNINGS AND PRECAUTIONS

- **Renal Impairment:** Evaluate the risks and benefits in patients with known renal impairment or taking nephrotoxic drugs; monitor renal function (5.1, 7.1, 8.6, 13.2)

- **Mesalamine-induced Acute Intolerance Syndrome:** Symptoms may be difficult to distinguish from an ulcerative colitis exacerbation; monitor for worsening symptoms; discontinue if acute intolerance syndrome suspected (5.2)
- **Hypersensitivity Reactions, including Myocarditis and Pericarditis:** Evaluate patients immediately and discontinue if a hypersensitivity reaction is suspected (5.3)
- **Hepatic Failure:** Evaluate the risks and benefits in patients with known liver impairment (5.4)

ADVERSE REACTIONS

The most common adverse reactions ($\geq 2\%$) are headache, nausea, nasopharyngitis, abdominal pain, and worsening of ulcerative colitis (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Warner Chilcott at 1-800-521-8813 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

DRUG INTERACTIONS

- **Nephrotoxic Agents including NSAIDs:** Increased risk of nephrotoxicity; monitor for changes in renal function and mesalamine-related adverse reactions. (7.1)
- **Azathioprine or 6-Mercaptopurine:** Increased risk of blood disorders; monitor complete blood cell counts and platelet counts (7.2)

USE IN SPECIFIC POPULATIONS

Geriatric Patients: Increased risk of blood dyscrasias; monitor complete blood cell counts and platelet counts (8.5)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 05/2016

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Administration Instructions
- 2.2 Dosage Information

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

- 5.1 Renal Impairment
- 5.2 Mesalamine-Induced Acute Intolerance Syndrome
- 5.3 Hypersensitivity Reactions
- 5.4 Hepatic Failure

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Nephrotoxic Agents, Including Non-Steroidal Anti-Inflammatory Drugs
- 7.2 Azathioprine or 6-Mercaptopurine

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

14.1 Moderately Active Ulcerative Colitis

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Asacol® HD is indicated for the treatment of moderately active ulcerative colitis in adults.

Limitations of Use:

Safety and effectiveness of Asacol HD beyond 6 weeks have not been established.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

- Do not substitute one Asacol HD 800 tablet for two mesalamine delayed-release 400 mg oral products [*see Clinical Pharmacology (12.3)*].
- Evaluate renal function prior to initiation of Asacol HD.
- Take Asacol HD tablets on an empty stomach, at least 1 hour before and 2 hours after a meal [*see Clinical Pharmacology (12.3)*].
- Swallow Asacol HD tablets whole. Do not cut, break or chew the tablets.
- Intact, partially intact, and/or tablet shells have been reported in the stool; Instruct patients to contact their physician if this occurs repeatedly.
- Protect Asacol HD tablets from moisture. Close the container tightly and leave desiccant pouches in the bottle along with the tablets.

2.2 Dosage Information

For the treatment of moderately active ulcerative colitis, the recommended dosage of Asacol HD in adults is 1600 mg (two 800 mg tablets) three times daily (total daily dosage of 4.8 grams) for a duration of 6 weeks.

3 DOSAGE FORMS AND STRENGTHS

Asacol HD delayed-release tablets: 800 mg (red-brown, capsule-shaped and imprinted with “WC 800” in black).

4 CONTRAINDICATIONS

Asacol HD is contraindicated in patients with known or suspected hypersensitivity to salicylates or aminosaliculates or to any of the ingredients of Asacol HD [*see Warnings and Precautions (5.3), Adverse Reactions (6.2), and Description (11)*].

5 WARNINGS AND PRECAUTIONS

5.1 Renal Impairment

Renal impairment, including minimal change nephropathy, acute and chronic interstitial nephritis, and, rarely, renal failure, has been reported in patients taking products such as Asacol HD that contain or are converted to mesalamine [*see Adverse Reactions (6.2)*].

Evaluate renal function prior to initiation of Asacol HD and periodically while on therapy. Evaluate the risks and benefits of using Asacol HD in patients with known renal impairment or history of renal disease or taking concomitant nephrotoxic drugs [*see Drug Interactions (7.1), Use in Specific Populations (8.6) and Nonclinical Toxicology (13.2)*].

5.2 Mesalamine-Induced Acute Intolerance Syndrome

Mesalamine has been associated with an acute intolerance syndrome that may be difficult to distinguish from an exacerbation of ulcerative colitis. Exacerbation of the symptoms of colitis has been reported in 2.3% of Asacol HD-treated patients in controlled clinical trials. This acute reaction, characterized by cramping, abdominal pain, bloody diarrhea, and occasionally by fever, headache, malaise, pruritus, rash, and conjunctivitis, has been reported after the initiation of Asacol HD tablets as well as other mesalamine products. Symptoms usually abate when Asacol HD tablets are discontinued.

5.3 Hypersensitivity Reactions

Hypersensitivity reactions have been reported in patients taking sulfasalazine. Some patients may have a similar reaction to Asacol HD tablets or to other compounds that contain or are converted to mesalamine.

As with sulfasalazine, mesalamine-induced hypersensitivity reactions may present as internal organ involvement, including myocarditis, pericarditis, nephritis, hepatitis, pneumonitis, and hematologic abnormalities. Evaluate patients immediately if signs or symptoms of a hypersensitivity reaction are present. Discontinue Asacol HD if an alternative etiology for the signs or symptoms cannot be established.

5.4 Hepatic Failure

There have been reports of hepatic failure in patients with pre-existing liver disease who have been administered mesalamine. Caution should be exercised when administering Asacol HD to patients with liver impairment.

6 ADVERSE REACTIONS

The most serious adverse reactions seen in Asacol HD clinical trials or with other products that contain mesalamine or are metabolized to mesalamine were:

- Renal Impairment [*see Warnings and Precautions (5.1)*]
- Mesalamine-Induced Acute Intolerance Syndrome [*see Warnings and Precautions (5.2)*]
- Hypersensitivity Reactions [*see Warnings and Precautions (5.3)*]
- Hepatic Failure [*see Warnings and Precautions (5.4)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Asacol HD has been evaluated in 896 patients with ulcerative colitis in controlled studies. Three six-week, active-controlled studies were conducted comparing Asacol HD 4.8 grams per day with mesalamine-delayed release tablets 2.4 grams per day in patients with mildly to moderately active ulcerative colitis. In these studies, 727 patients were dosed with Asacol HD tablets and 732 patients were dosed with mesalamine delayed-release tablets.

The most common reactions reported in the Asacol HD group were headache (4.7%), nausea (2.8%), nasopharyngitis (2.5%), abdominal pain (2.3%), diarrhea (1.7%), and dyspepsia (1.7%); Table 1 enumerates adverse reactions that occurred in the three studies. The most common reactions in patients with moderately active ulcerative colitis (602 patients dosed with Asacol HD and 618 patients dosed with mesalamine delayed-release 400 mg) were the same as all treated patients.

Discontinuations due to adverse reactions occurred in 3.9% of patients in the Asacol HD group and in 4.2% of patients in the mesalamine delayed-release tablet comparator group. The most common cause for discontinuation was gastrointestinal symptoms associated with ulcerative colitis.

Serious adverse reactions occurred in 0.8% of patients in the Asacol HD group and in 1.8% of patients in the mesalamine delayed-release tablet comparator group. The majority involved the gastrointestinal system.

Table 1. Adverse Reactions Occurring in \geq1% of All Treated Patients (Three studies combined)		
Adverse Reaction	Mesalamine delayed-release 2.4 grams per day (400 mg Tablet) (N = 732)	Asacol HD 4.8 grams per day (800 mg Tablet) (N = 727)
Headache	4.9 %	4.7 %
Nausea	2.9 %	2.8 %
Nasopharyngitis	1.4 %	2.5 %
Abdominal pain	2.3 %	2.3 %
Diarrhea	1.9 %	1.7 %
Dyspepsia	0.8 %	1.7 %
Vomiting	1.6 %	1.4 %
Flatulence	0.7 %	1.2 %
Influenza	1.2 %	1.0 %
Pyrexia	1.2 %	0.7 %
Cough	1.4 %	0.3 %
N = number of patients within specified treatment group Percent = percentage of patients in category and treatment group		

6.2 Postmarketing Experience

In addition to the adverse reactions reported above in clinical trials involving the Asacol HD tablet, the adverse events listed below have been reported in postmarketing experience with other mesalamine-containing products or products that are metabolized to mesalamine. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a Whole: Facial edema, edema, peripheral edema, asthenia, chills, infection, malaise, pain, neck pain, chest pain, back pain, abdominal enlargement, lupus-like syndrome, drug fever (rare).

Cardiovascular: Pericarditis (rare) and myocarditis (rare) [*see Warnings and Precautions (5.3)*], pericardial effusion, vasodilation, migraine.

Gastrointestinal: Dry mouth, stomatitis, oral ulcers, anorexia, increased appetite, eructation, pancreatitis, cholecystitis, gastritis, gastroenteritis, gastrointestinal bleeding, perforated peptic ulcer (rare), constipation, hemorrhoids, rectal hemorrhage, bloody diarrhea, tenesmus, stool abnormality.

Hepatic: There have been rare reports of hepatotoxicity, including jaundice, cholestatic jaundice, hepatitis, and possible hepatocellular damage including liver necrosis and liver failure. Some of these cases were fatal. Asymptomatic elevations of liver enzymes which usually resolve during continued use or with discontinuation of the drug have also been reported. One case of Kawasaki-like syndrome, that included changes in liver enzymes, was also reported [*see Warnings and Precautions (5.4)*].

Hematologic: Agranulocytosis (rare), aplastic anemia (rare), anemia, thrombocytopenia, leukopenia, eosinophilia, lymphadenopathy.

Musculoskeletal: Gout, rheumatoid arthritis, arthritis, arthralgia, joint disorder, myalgia, hypertonia.

Neurological/Psychiatric: Anxiety, depression, somnolence, insomnia, nervousness, confusion, emotional lability, dizziness, vertigo, tremor, paresthesia, hyperesthesia, peripheral neuropathy (rare),

Guillain-Barré syndrome (rare), and transverse myelitis (rare).

Respiratory/Pulmonary: Sinusitis, rhinitis, pharyngitis, asthma exacerbation, pleuritis, bronchitis, eosinophilic pneumonia, interstitial pneumonitis.

Skin: Alopecia, psoriasis (rare), pyoderma gangrenosum (rare), erythema nodosum, acne, dry skin, sweating, pruritus, urticaria, rash.

Special Senses: Ear pain, tinnitus, ear congestion, ear disorder, conjunctivitis, eye pain, blurred vision, vision abnormality, taste perversion.

Renal/Urogenital: Renal failure (rare), interstitial nephritis, minimal change nephropathy [*see Warnings and Precautions (5.1)*], dysuria, urinary frequency and urgency, hematuria, epididymitis, decreased libido, dysmenorrhea, menorrhagia.

Laboratory Abnormalities: Elevated AST (SGOT) or ALT (SGPT), elevated alkaline phosphatase, elevated GGT, elevated LDH, elevated bilirubin, elevated serum creatinine and BUN.

7 DRUG INTERACTIONS

7.1 Nephrotoxic Agents, Including Non-Steroidal Anti-Inflammatory Drugs

The concurrent use of mesalamine with known nephrotoxic agents, including nonsteroidal anti-inflammatory drugs (NSAIDs) may increase the risk of nephrotoxicity. Monitor

patients taking nephrotoxic drugs for changes in renal function and mesalamine-related adverse reactions [see *Warnings and Precautions (5.1)*].

7.2 Azathioprine or 6-Mercaptopurine

The concurrent use of mesalamine with azathioprine or 6-mercaptopurine may increase the risk for blood disorders. If concomitant use of Asacol HD and azathioprine or 6-mercaptopurine cannot be avoided, monitor blood tests, including complete blood cell counts and platelet counts.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Limited published data on mesalamine use in pregnant women are insufficient to inform a drug-associated risk. No fetal harm was observed in animal reproduction studies of mesalamine in rats and rabbits at oral doses approximately 0.97 times (rat) and 1.95 times (rabbit) the recommended human dose [see *Data*].

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

Reproduction studies with mesalamine were performed during organogenesis in rats and rabbits at oral doses up to 480 mg/kg/day. There was no evidence of harm to the fetus. These mesalamine doses were about 0.97 times (rat) and 1.95 times (rabbit) the recommended human dose of 4.8 grams per day, based on body surface area.

8.2 Lactation

Risk Summary

Mesalamine and its N-acetyl metabolite are present in human milk in undetectable to small amounts [see *Data*]. There are limited reports of diarrhea in breastfed infants. There is no information on the effects of the drug on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Asacol HD and any potential adverse effects on the breastfed infant from the drug or from the underlying maternal condition.

Clinical Considerations

Monitor breastfed infants for diarrhea.

Data

Human Data

In published lactation studies, maternal mesalamine doses from various oral and rectal formulations and products ranged from 500 mg to 3 g daily. The concentration of mesalamine in milk ranged from non-detectable to 0.11 mg/L. The concentration of the N-acetyl-5-aminosalicylic acid metabolite ranged from 5 to 18.1 mg/L. Based on these concentrations, estimated infant daily dosages for an exclusively breastfed infant are 0 to 0.017 mg/kg/day of mesalamine and 0.75 to 2.72 mg/kg/day of N-acetyl-5-aminosalicylic acid.

8.4 Pediatric Use

Safety and effectiveness of Asacol HD in pediatric patients have not been established. See the prescribing information for other approved mesalamine products for the safety and effectiveness of these products in pediatric patients.

8.5 Geriatric Use

Clinical studies of Asacol HD did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently than younger patients. Reports from uncontrolled clinical studies and postmarketing experience for another oral formulation of mesalamine suggest a higher incidence of blood dyscrasias (agranulocytosis, neutropenia, pancytopenia) in patients who were 65 years or older compared to younger patients. Monitor complete blood cell counts and platelet counts in elderly patients during therapy with Asacol HD.

In general, the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy in elderly patients should be considered when prescribing Asacol HD [*see Use in Specific Populations (8.6)*].

8.6 Renal Impairment

Mesalamine is known to be substantially excreted by the kidney, and the risk of adverse reactions may be greater in patients with impaired renal function. Evaluate renal function in all patients prior to initiation and periodically while on Asacol HD therapy. Monitor patients with known renal impairment or history of renal disease or taking nephrotoxic drugs for decreased renal function and mesalamine-related adverse reactions [*see Warnings and Precautions (5.1), Drug Interactions (7.1) and Adverse Reactions (6.2)*].


10 OVERDOSAGE

There is no specific antidote for mesalamine overdose and treatment for suspected acute severe toxicity with Asacol HD should be symptomatic and supportive. This may include prevention of further gastrointestinal tract absorption, correction of fluid electrolyte imbalance, and maintenance of adequate renal function. Asacol HD is a pH dependent delayed-release product and this factor should be considered when treating a suspected overdose.

Single oral doses of 5000 mg/kg mesalamine suspension in mice (approximately 4.2 times the recommended human dose of Asacol HD based on body surface area), 4595 mg/kg in rats (approximately 7.8 times the recommended human dose of Asacol HD based on body surface area) and 3000 mg/kg in cynomolgus monkeys (approximately 10 times the recommended human dose of Asacol HD based on body surface area) were lethal.

11 DESCRIPTION

Each Asacol HD delayed-release tablet for oral administration contains 800 mg of mesalamine, an aminosalicylate. Asacol HD delayed-release tablets have an outer protective coat consisting of a combination of acrylic based resins, Eudragit S (methacrylic acid and methyl methacrylate copolymer (1:2), NF) and Eudragit L (methacrylic acid and methyl methacrylate copolymer (1:1), NF). The inner coat consists of an acrylic based resin, Eudragit S, which dissolves at pH 7 or greater, releasing mesalamine in the terminal ileum and beyond for topical anti-inflammatory action in the colon. Mesalamine (also referred to as 5-aminosalicylic acid or 5-ASA) has the chemical name 5-amino-2-hydroxybenzoic acid; its structural formula is:



Molecular Weight: 153.1
Molecular Formula: C₇H₇NO₃

Inactive Ingredients: Each tablet contains colloidal silicon dioxide, dibutyl sebacate, edible black ink, ferric oxide red, ferric oxide yellow, lactose monohydrate, magnesium stearate, methacrylic acid and methyl methacrylate copolymer (1:2) (Eudragit S), methacrylic acid and methyl methacrylate copolymer (1:1) (Eudragit L), polyethylene glycol, povidone, sodium starch glycolate, and talc.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of mesalamine is unknown, but appears to be topical rather than systemic. Mucosal production of arachidonic acid metabolites, both through the cyclooxygenase pathways, that is, prostanoids, and through the lipoxygenase pathways, that is, leukotrienes and hydroxyeicosatetraenoic acids, is increased in patients with chronic inflammatory bowel disease, and it is possible that mesalamine diminishes inflammation by blocking cyclooxygenase and inhibiting prostaglandin production in the colon.

12.3 Pharmacokinetics

Absorption

Plasma concentrations of mesalamine (5-aminosalicylic acid; 5-ASA) and its metabolite, N-acetyl-5-aminosalicylic acid (N-Ac-5-ASA) are highly variable following administration of Asacol HD tablets. Following single dose oral administration of Asacol HD 800 mg tablet in healthy subjects (N = 139) under fasted conditions, the mean C_{max} , AUC_{8-48h} and $AUC_{0-t_{1/2c}}$ values were 208 ng/mL, 2296 ng.h/mL, and 2533 ng.h/mL, respectively. The median [range] T_{max} for mesalamine following administration of Asacol HD 800 mg tablet was approximately 24 hours [4 to 72 hours], reflecting the delayed-release characteristics of the formulation.

Based on cumulative urinary recovery of mesalamine and N-Ac-5-ASA from single dose studies in healthy subjects, approximately 20% of the orally administered mesalamine in Asacol HD tablets is systemically absorbed.

Food Effect: A high calorie (800 to 1000 calories), high fat (approximately 50 % of total caloric content) meal increased mesalamine C_{max} by 2.4-fold and mesalamine systemic exposure (AUC_{8-48} and $AUC_{0-t_{1/2c}}$) by 2.8-fold; the median lag-time increased by 8 hours and median t_{max} by 6 hours (from 24 to 30 hours) [see *Dosage and Administration (2.1)*].

Comparative exposure between one Asacol HD 800 mg tablet and two mesalamine delayed-release 400 mg oral products is unknown [see *Dosage and Administration (2.1)*].

Elimination

Metabolism

The absorbed mesalamine is acetylated in the gut mucosal wall and by the liver to N-Ac-5-ASA.

Excretion

Absorbed mesalamine is excreted mainly by the kidneys as N-acetyl-5-aminosalicylic acid. Unabsorbed mesalamine is excreted in feces.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Dietary mesalamine was not carcinogenic in rats at doses as high as 480 mg/kg/day, or in mice at 2000 mg/kg/day. These doses are approximately 0.97 and 2.0 times the 4.8 grams per day Asacol HD dose (based on body surface area). Mesalamine was not genotoxic in the Ames test, the Chinese hamster ovary cell chromosomal aberration assay, and the mouse micronucleus test. Mesalamine, at oral doses up to 480 mg/kg/day (about 0.97 times the recommended human treatment dose based on body surface area), was found to have no effect on fertility or reproductive performance of male and female rats.

13.2 Animal Toxicology and/or Pharmacology

In animal studies (rats, mice, dogs), the kidney was the principal organ for toxicity. (In the following, comparisons of animal dosing to recommended human dosing are based on body surface area and a 4.8 grams per day dose for a 60 kg person.)

Mesalamine causes renal papillary necrosis in rats at single doses of approximately 750 mg/kg to 1000 mg/kg (1.5 to 2.0 times the recommended human dose). Doses of 170 and 360 mg/kg/day (about 0.3 and 0.73 times the recommended human dose) given to rats for six months produced papillary necrosis, papillary edema, tubular degeneration, tubular mineralization, and urothelial hyperplasia.

In mice, oral doses of 4000 mg/kg/day (approximately 4.1 times the recommended human dose) for three months produced tubular nephrosis, multifocal/diffuse tubulo-interstitial inflammation, and multifocal/diffuse papillary necrosis.

In dogs, single doses of 6000 mg (approximately 6.25 times the recommended human dose) of delayed-release mesalamine tablets resulted in renal papillary necrosis but were not fatal. Renal changes have occurred in dogs given chronic administration of mesalamine at doses of 80 mg/kg/day (0.5 times the recommended human dose).

14 CLINICAL STUDIES

14.1 Moderately Active Ulcerative Colitis

The efficacy of Asacol HD at 4.8 grams per day was studied in a six-week, randomized, double-blind, active-controlled study in 772 patients with moderately active ulcerative colitis (UC). Moderately active UC was defined as a Physician's Global Assessment (PGA) score of 2; the PGA is a four-point scale (0 to 3) that encompasses the clinical assessments of rectal bleeding, stool frequency, and sigmoidoscopy findings.

Patients were randomized 1:1 to the Asacol HD 4.8 grams per day group (two Asacol HD tablets three times a day) or the mesalamine delayed-release 2.4 grams per day group (two mesalamine delayed-release 400 mg tablets three times a day).

Patients characteristically had a history of previous use of oral 5-ASAs (86%), steroids (41%), and rectal therapies (49%), and demonstrated clinical symptoms of three or more stools over normal per day (87%) and obvious blood in the stool most or all of the time (70%). The study population was primarily Caucasian (97%), had a mean age of 43 years (8% aged 65 years or older), and included slightly more males (56%) than females (44%).

The primary endpoint was treatment success defined as improvement from baseline to Week 6 based on the PGA. Treatment success rates were similar in the two groups: 70% in the Asacol HD group and 66% in the Asacol group (difference: 5%; 95% CI: [-1.9%, 11.2%]).

A second controlled study supported the efficacy of Asacol HD at 4.8 grams per day. Treatment success was 72% in patients with moderately active UC treated with Asacol HD.

16 HOW SUPPLIED/STORAGE AND HANDLING

Asacol[®] HD (mesalamine) delayed-release tablets are available as red-brown, capsule-shaped tablets containing 800 mg mesalamine and imprinted with “WC 800” in black.

NDC 0430-0784-27 Bottle of 180 tablets

Store and dispense in the original bottle, protect from moisture, and keep the bottle tightly closed. Do not remove desiccant pouch (silica gel) from bottle.

Store at controlled room temperature, 20° to 25° C (68° to 77° F); excursions are permitted 15° to 30° C (59° to 86° F). [See USP Controlled Room Temperature]

17 PATIENT COUNSELING INFORMATION

Administration [see Dosage and Administration (2.1)]

- Inform patients that if they are switching from a previous oral mesalamine therapy to Asacol HD to discontinue their previous oral mesalamine therapy and follow the dosing instructions for Asacol HD. One Asacol HD 800 mg tablet is not substitutable for two mesalamine delayed-release 400 mg oral products.
- Inform patients to take Asacol HD tablets on an empty stomach, at least 1 hour before and 2 hours after a meal.
- Instruct patients to swallow the Asacol HD tablets whole, taking care not to break, cut, or chew the tablets, because the coating is an important part of the delayed-release formulation.

- Inform patients that intact, partially intact, and/or tablet shells have been reported in the stool. Instruct patients to contact their physician if this occurs repeatedly.
- Instruct patients to protect Asacol HD tablets from moisture. Instruct patients to close the container tightly and to leave desiccant pouches in the bottle along with the tablets.

Renal Impairment

- Inform patients that Asacol HD may decrease their renal function, especially if they have known renal impairment or are taking nephrotoxic drugs, and periodic monitoring of renal function will be performed while they are on therapy. Advise patients to complete all blood tests ordered by their physician [*see Warnings and Precautions (5.1)*].

Mesalamine-Induced Acute Intolerance Syndrome

- Instruct patients to report to their physician if they experience new or worsening symptoms of cramping, abdominal pain, bloody diarrhea, and sometimes fever, headache, and rash [*see Warnings and Precautions (5.2)*].

Hypersensitivity Reactions

- Inform patients of the signs and symptoms of hypersensitivity reactions, and advise them seek immediate medical care should signs and symptoms occur [*see Warnings and Precautions (5.3)*].

Hepatic Failure

- Inform patients with known liver disease of the signs and symptoms of worsening liver function and advise them to report to their physician if they experience such signs or symptoms [*see Warnings and Precautions (5.4)*].

Blood Disorders

- Inform elderly patients and those taking azathioprine or 6-mercaptopurine of the risk for blood disorders and the need for periodic monitoring of complete blood cell counts and platelet counts while on therapy. Advise patients to complete all blood tests ordered by their physician [*see Drug Interactions (7.2), Use in Specific Populations (8.5)*].

Manufactured by:
Warner Chilcott
Deutschland GmbH D-

64331 Weiterstadt,
Germany

Marketed by:
Warner Chilcott
(US), LLC
Rockaway, NJ
07866

Under license from Medeva Pharma Suisse AG (registered trademark owner). U.S. Patent

No. 6,893,662.



Dispense in original container. Store at controlled room temperature 20° - 25° C (68° - 77° F) [See USP].

For the treatment of moderately active ulcerative colitis, the recommended dose of Asacol HD in adults is 1600 mg (two 800 mg tablets) three times daily (total daily dosage of 4.8 grams) for a duration of 6 weeks.

See Package Outsert for Full Prescribing Information.

Visit us at our website: www.asacolHD.com

NDC 0430-0783-95
Rx Only

SAMPLE-Not for Sale

Asacol[®] HD
(mesalamine)
delayed-release tablets
800 mg per tablet

Do not substitute one Asacol HD 800 mg tablet for two mesalamine delayed-release 400 mg oral products

EXP
LOT

12 Tablets
XXXXXXX

Swallow Asacol HD tablets whole. Do not cut, break, or chew the tablets.


Mfg. by: Warner Chilcott Deutschland GmbH
D-64331 Weiterstadt, Germany
Mkt. by: Warner Chilcott (US), LLC, Rockaway, NJ 07866
1-800-521-8813
Under license from Medeva Suisse Pharma AG (registered trademark owner).




Global Labeling and Packaging Specifications

Part number TBD (draft 4/12/16)
Drawing number DL107

- Artwork is actual size
- Drop template/die line and notes before processing






Global Labeling and Packaging Specifications

Part number TBD (draft 4/12/16)
 Drawing number DL109

- Artwork is actual size
 - Drop template/die line and notes before processing



NDC 0430-0784-27

Asacol[®] HD (mesalamine) delayed-release tablets

800 mg per tablet

Do not substitute one Asacol HD 800 mg tablet for two mesalamine delayed-release 400 mg oral products



180 Tablets Rx Only
asacolHD.com


Dispense in original container. Store at controlled room temperature 20° - 25° C (68° - 77° F) [See USP].

For the treatment of moderately active ulcerative colitis, the recommended dose of Asacol HD in adults is 1600 mg (two 800 mg tablets) three times daily (total daily dosage of 4.8 grams) for a duration of 6 weeks.

See Package Outsert for Full Prescribing Information.

Swallow Asacol HD tablets whole. Do not cut, break, or chew the tablets.

Mfg. by: Warner Chilcott Deutschland GmbH
D-64331 Weiterstadt, Germany
Mkt. by: Warner Chilcott (US), LLC
Rockaway, NJ 07866
1-800-521-8813
Under license from Medeva Suisse Pharma AG (registered trademark owner).



XXXXXXXX

EXP
LOT


XXXXXXXX



Global Labeling and Packaging Specifications

Part number TBD (draft 4/12/16)
Drawing number DL108

- Artwork is actual size
- Drop template/die line and notes before processing



This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOYCE A KORVICK
05/05/2016