

FACT SHEET FOR INDIVIDUALS

OraSure Technologies, Inc.

InteliSwab™ COVID-19 Rapid Test

Updated: January 27, 2022

Coronavirus
Disease 2019
(COVID-19)

You are provided this Fact Sheet because you obtained the InteliSwab™ COVID-19 Rapid Test for testing yourself or dependants for the proteins from the virus that causes COVID-19. The intended use of this test is for testing individuals within 7 days of COVID-19 symptom onset or individuals with or without symptoms or other epidemiological reasons to suspect COVID-19 when tested twice over three days with at least 24 hours (and no more than 48 hours) between tests.

This Fact Sheet contains information to help you understand the risks and benefits of using this test for the detection of proteins from the virus that causes COVID-19. After reading this Fact Sheet, if you have questions or would like to discuss the information provided, please talk to your healthcare provider.

For the most up to date information on COVID-19 please visit the CDC Coronavirus Disease 2019 (COVID-19) webpage:

<https://www.cdc.gov/COVID19>

What is COVID-19?

COVID-19 is caused by the SARS-CoV-2 virus which is a new virus in humans causing a contagious respiratory illness. COVID-19 can present with a mild to severe illness, although some people infected with COVID-19 may have no symptoms at all. Older adults and people of any age who have underlying medical conditions have a higher risk of severe illness from COVID-19. Serious outcomes of COVID-19 include hospitalization and death. The SARS-CoV-2 virus can be spread to others not just while one is sick, but even before a person shows signs or symptoms of being sick (e.g., fever, coughing, difficulty breathing, etc.). A full list of symptoms of COVID-19 can be found at the following link: <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>.

What is the InteliSwab™ COVID-19 Rapid Test?

The InteliSwab™ COVID-19 Rapid Test is a type of test called an antigen test. Antigen tests are designed to detect proteins from the virus that causes COVID-19 in anterior nasal swab specimens. The InteliSwab™ COVID-19 Rapid Test is intended for the qualitative detection of the nucleocapsid protein antigen from SARS-CoV-2 from individuals within 7 days of COVID-19 symptom onset or individuals with or without symptoms or other epidemiological reasons to suspect COVID-19 when tested twice over three days with at least 24 hours (and no more than 48 hours) between tests.

What are the known and potential risks and benefits of the test?

Potential risks include:

- Possible discomfort or other complications that can happen during sample collection.
- Possible incorrect test result (see below for more information).

Potential benefits include:

- The results, along with other information, can help your healthcare provider make informed recommendations about your care.
- The results of this test may help limit the spread of COVID-19 to your family and those you come in contact with.

What does it mean if I have a positive test result?

If you have a positive test result, it is very likely that you have COVID-19 because proteins from the virus that causes COVID-19 were found in your sample. Therefore, it is also likely that you may be placed in isolation to avoid spreading the virus to others. There is a very small chance that this test can give a positive result that is wrong (a false positive result). If you test positive with the InteliSwab™ COVID-19 Rapid Test you should self-isolate and seek follow-up care with your healthcare provider as additional testing may be necessary. Your healthcare provider will work with you to determine how best to care for you based on your test

Where can I go for updates and more information? The most up-to-date information on COVID-19 is available at the CDC General webpage: <https://www.cdc.gov/COVID19>. In addition, please also contact your healthcare provider with any questions/concerns.

FACT SHEET FOR INDIVIDUALS

OraSure Technologies, Inc.

InteliSwab™ COVID-19 Rapid Test

Updated: January 27, 2022

Coronavirus
Disease 2019
(COVID-19)

result(s) along with other factors of your medical history, and your symptoms.

What does it mean if I have a negative test result?

A negative test result means that proteins from the virus that causes COVID-19 were not found in your sample. It is possible for this test to give a negative result that is incorrect (false negative) in some people with COVID-19. This means that you could possibly still have COVID-19 even though the test is negative. The amount of antigen in a sample may decrease the longer you have symptoms of infection. In symptomatic people, specimens collected after you have had symptoms for more than seven days may be more likely to be negative compared to a molecular assay.

If you test negative and continue to experience COVID-19 like symptoms of fever, cough and/or shortness of breath you should seek follow up care with your healthcare provider. For example, your healthcare provider may suggest you need another test to determine if you have contracted the virus causing COVID-19. If you are concerned about your COVID-19 infection status after testing or think you may need follow up testing, please contact your healthcare provider.

The performance of this test was established based on the evaluation of a limited number of clinical specimens. The clinical performance has not been established in all circulating variants but is anticipated to be reflective of the prevalent variants in circulation at the time and location of the clinical evaluation. Performance at the time of testing may vary depending on the variants circulating, including newly emerging strains of SARS-CoV-2 and their prevalence, which change over time.

What is serial testing?

Serial testing is when a single person is tested for COVID-19 more than once. Because antigen tests are less sensitive than other COVID-19 tests and false results may occur, repeated testing may identify more individuals with COVID-19 than a single test. By

repeating testing, it may be possible to more quickly identify cases of COVID-19 and reduce spread of infection. Additional testing with molecular COVID-19 test may be necessary, depending on your individual risk factors and test results.

It is important that you work with your healthcare provider to help you understand the next steps you should take

What are the differences between antigen tests and other COVID-19 tests?

There are different kinds of tests for diagnosing COVID-19. Molecular tests (also known as PCR tests) detect genetic material from the virus. Antigen tests detect proteins from the virus. Antigen tests are very specific for the virus, but are not as sensitive as molecular tests. This means that a positive result is highly accurate, but a negative result does not rule out infection.

If your test result is negative, you should discuss with your healthcare provider whether an additional molecular test would help with your care, and when you should discontinue home isolation. If you do not have an additional test to determine if you are infected and may spread the infection to others, the CDC currently recommends that you should stay home until three things have happened:

- You have had no fever for at least 24 hours (that is three full days of no fever without the use of medicine that reduces fevers).

AND

- Other symptoms of COVID-19 have improved (for example, when your cough or shortness of breath has improved). Loss of taste and smell may persist for weeks or months after recovery and need not delay the end of isolation.

AND

- At least 10 days have passed since your symptoms first appeared.

Where can I go for updates and more information? The most up-to-date information on COVID-19 is available at the CDC General webpage: <https://www.cdc.gov/COVID19>. In addition, please also contact your healthcare provider with any questions/concerns.

FACT SHEET FOR INDIVIDUALS

OraSure Technologies, Inc.

InteliSwab™ COVID-19 Rapid Test

Updated: January 27, 2022

Coronavirus
Disease 2019
(COVID-19)

For more information, the CDC has provided guidelines on how to prevent the spread of COVID-19 if you are sick: <https://www.cdc.gov/coronavirus/2019-ncov/downloads/sick-with-2019-nCoV-fact-sheet.pdf>

Is this test FDA-approved or cleared?

No. This test is not yet approved or cleared by the United States FDA. FDA may issue an Emergency Use Authorization (EUA) when certain criteria are met, which includes that there are no adequate, approved, available alternatives. The EUA for this test is supported by the Secretary of Health and Human Service's (HHS's) declaration that circumstances exist to justify the emergency use of in vitro diagnostics for the detection and/or diagnosis of the virus that causes COVID-19. This EUA will remain in effect (meaning this test can be used) for the duration of the COVID-19 declaration justifying emergency of IVDs, unless it is terminated or revoked by FDA (after which the test may no longer be used).

What are the approved alternatives?

There are no approved available alternative antigen tests. Any tests that have received full marketing status (e.g., cleared, approved), as opposed to an EUA, by FDA can be found by searching the medical device databases here: <https://www.fda.gov/medical-devices/device-advice-comprehensive-regulatory-assistance/medical-device-databases>. A cleared or approved test should be used instead of a test made available under an EUA, when appropriate and available. FDA has issued EUAs for other tests that can be found at: <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization>.

Where can I go for updates and more information? The most up-to-date information on COVID-19 is available at the CDC General webpage: <https://www.cdc.gov/COVID19>. In addition, please also contact your healthcare provider with any questions/concerns.
