

NOT APPROVED BY FDA – Legally marketed as an FDA Indexed Product under MIF 900-013.  
FOR USE IN FERRETS ONLY.  
Extra-label use is prohibited.  
This product is not to be used in animals intended for use as food for humans or other animals.

CAUTION: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.


# Suprelorin® F

## (DESLORELIN ACETATE) 4.7 mg Implant

### DESCRIPTION

**Suprelorin® F** (4.7 mg) Implant is a synthetic GnRH analogue (deslorelin acetate) in a biocompatible, slow release subcutaneous implant. The implant is a solid, opaque, white to pale yellow cylinder, 2.3 mm x 12.5 mm in length and weighing 50 mg. The **Suprelorin® F** (4.7 mg) Implant comes pre-loaded in an implanting needle. Each implant contains 4.7 mg deslorelin (as deslorelin acetate) in an inert matrix.

### Chemical Structure – Deslorelin Acetate



[(6-D-tryptophan-9-(N-ethyl-L-prolinamide)-10-deglycinamide]GnRH

### INDICATIONS

**Suprelorin® F** (4.7 mg) Implant is indicated for the management of adrenal gland cortical disease in the male and female domestic ferret.

### DOSAGE AND ADMINISTRATION

The recommended dosage is one, 4.7 mg implant per ferret every 12 months. Appropriate clinical monitoring is suggested to determine that the symptoms of adrenal gland disease are being adequately controlled.

Do not use if the foil pouch is damaged.

Remove the luer lock cap from the implanting needle. Attach the actuator syringe to the implanting needle using the luer lock connection. One implant should be implanted subcutaneously at the dorsal aspect of the base of the neck. Administer only one implant per ferret. Select the implant site by locating the area of the back midway between the shoulder blades. It is not necessary to prepare the implantation site. If the hair is long, a small section may be clipped if required. Lift the loose skin between the shoulder blades. Insert the entire length of the needle subcutaneously. Fully depress the actuator syringe plunger. Press the skin at the insertion site as the needle is slowly withdrawn, and maintain pressure for 30 seconds. Examine the implanting needle to verify that the implant has not remained within the needle, and that the blue plastic spacer is visible at the tip of the needle. It may be possible to palpate the implant *in situ*. The biocompatible implant does not require removal. Wash hands after use.

Repeat treatment every 12 months to maintain efficacy. Appropriate endocrine testing and clinical monitoring should be performed at appropriate intervals to monitor the response to therapy.

### CONTRAINDICATIONS

Do not use this product in ferrets with known hypersensitivity to deslorelin acetate or other synthetic hormones.

### HUMAN SAFETY WARNINGS

**KEEP OUT OF REACH OF CHILDREN.** DO NOT HANDLE THIS PRODUCT IF YOU ARE PREGNANT OR NURSING OR SUSPECT YOU MAY BE PREGNANT. Accidental administration may lead to a disruption of the menstrual cycle. Avoid direct skin contact with the implant; if skin contact occurs, wash the affected area immediately with soap and water. The use of gloves is advised. As with all injectable drugs causing profound physiological effects, routine precautions should be employed by practitioners when handling and using **Suprelorin® F** (4.7 mg) Implant to prevent accidental injection. In case of accidental human injection, a physician should be consulted and the implant should be removed.

### PRECAUTION

Do not use in animals intended for breeding. The safe use of this product has not been evaluated in pregnant or lactating ferrets.

### ADVERSE REACTIONS

It is possible that treated ferrets will exhibit signs of soreness and swelling at the implantation site which should resolve over one or two weeks. Undesirable histology at the site of implantation has not been reported in other species (canine). Other reported side effects include: weight gain, lethargy and failure to respond to therapy.

To report suspected adverse drug events, please call Virbac at 1-855-647-3747.

### PHARMACOLOGY

**Suprelorin® F** (4.7 mg) Implant is a controlled release implant containing the GnRH agonist deslorelin. Deslorelin acetate suppresses the reproductive endocrine system, preventing production of pituitary and gonadal hormones. Deslorelin acetate has not been shown to reduce the size of adrenal tumors and is not considered curative.

### DISPOSAL

Each implanting needle (sterile) is a single use device. Used needles should immediately be placed in a designated and appropriately labeled "sharps" container. Each actuator syringe (non-sterile) is a multi-use device and should be saved for future use with the remaining implant(s) in the carton. Unused implants should be disposed of in accordance with local environmental requirements.

### STORAGE

Store at temperatures between 2° and 8° C (36° and 46° F). Do not freeze.

### HOW SUPPLIED

Five (5) or two (2) implants pre-loaded in implanting needles and individually packaged per carton.

For technical assistance, to request an MSDS, or to report suspected adverse drug events, please call Virbac at 1-855-647-3747.



Manufactured for:  
Virbac AH, Inc.  
Fort Worth, Texas

Product of Australia

MIF 900-013

Revision 06/2012  
L-2000-F-US-1

Suprelorin® F

**INDICATION:** For the Management of Adrenal Gland Cortical Disease in the Male and Female Domestic Ferret.

**DIRECTIONS FOR USE:** Refer to the package insert for complete directions for use. Each implant contains 4.7 mg of deslorelin acetate. Do not use if the foil pouch is damaged. Administer one implant per ferret. One implant should be injected subcutaneously at the dorsal base of the neck.


**USER SAFETY:** Avoid direct skin contact with the implant. DO NOT HANDLE OR ADMINISTER THIS PRODUCT IF YOU ARE PREGNANT OR NURSING OR SUSPECT YOU MAY BE PREGNANT.

**WARNINGS:** KEEP OUT OF THE REACH OF CHILDREN. For use in ferrets only. Do not use in animals intended for breeding.

For technical assistance, to request an MSDS, or to report suspected adverse drug events, please call Virbac at 1-855-647-3747.

**DISPOSAL:** The implanting needle is a single use device and should be disposed of after use in the appropriate receptacle for sharp medical waste. The actuator syringe can be re-used. Unused implants should be disposed of in accordance with local environmental requirements.

**STORAGE:** Store at temperatures between 2° and 8°C (36° and 46°F). Do not freeze.



0 84942 44402 3

**Virbac**  
ANIMAL HEALTH

NOT APPROVED BY FDA - Legally marketed as an FDA Indexed Product under MIF 900-013.

FOR USE IN FERRETS ONLY. Extra-label use is prohibited. This product is not to be used in animals intended for use as food for humans or other animals.

# Suprelorin® F

(DESLORELIN ACETATE)  
4.7mg IMPLANT

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian

Box contains 2 implants pre-loaded in sterile implanting needles and 1 non-sterile actuator syringe



Product of Australia

Distributed in the U.S.A. by:  
Virbac AH, Inc.  
3200 Meacham Boulevard  
Fort Worth, Texas 76137  
1-855-647-3747

C-2002-F-US-1

LOT:  
EXP:

44402

**INDICATION:** For the Management of Adrenal Gland Cortical Disease in the Male and Female Domestic Ferret.

**DIRECTIONS FOR USE:** Refer to the package insert for complete directions for use. Each implant contains 4.7 mg of deslorelin acetate. Do not use if the foil pouch is damaged. Administer one implant per ferret. One implant should be injected subcutaneously at the dorsal base of the neck.

**USER SAFETY:** Avoid direct skin contact with the implant. DO NOT HANDLE OR ADMINISTER THIS PRODUCT IF YOU ARE PREGNANT OR NURSING OR SUSPECT YOU MAY BE PREGNANT.

**WARNINGS:** KEEP OUT OF THE REACH OF CHILDREN. For use in ferrets only. Do not use in animals intended for breeding.

For technical assistance, to request an MSDS, or to report suspected adverse drug events, please call Virbac at 1-855-647-3747.

**DISPOSAL:** The implanting needle is a single use device and should be disposed of after use in the appropriate receptacle for sharp medical waste. The actuator syringe can be re-used. Unused implants should be disposed of in accordance with local environmental requirements.

**STORAGE:** Store at temperatures between 2° and 8°C (36° and 46°F). Do not freeze.



0 84942 44405 4


**Virbac**  
ANIMAL HEALTH

NOT APPROVED BY FDA - Legally marketed as an FDA  
Indexed Product under MIF 900-013.  
FOR USE IN FERRETS ONLY. Extra-label use is prohibited.  
This product is not to be used in animals intended for  
use as food for humans or other animals.

**Suprelorin® F**  
**(DESLORELIN ACETATE)**  
**4.7mg IMPLANT**

**CAUTION:** Federal law  
restricts this drug to use by  
or on the order of a  
licensed veterinarian.

Box contains 5 implants pre-loaded in sterile implanting  
needles and 1 non-sterile actuator syringe



Lot :  
Exp :

Product of Australia

Distributed in the U.S.A. by:  
Virbac AH, Inc.  
3200 Meacham Boulevard  
Fort Worth, Texas 76137  
1-855-647-3747

44405

C-2005-F-US-1

**DIRECTIONS FOR USE:** Refer to package insert for complete directions for use.

**WARNING:** KEEP OUT OF THE REACH OF CHILDREN

**STORAGE:** Store at temperatures between 2° and 8°C (36° and 46°F). Do not freeze.

For technical assistance, to request an MSDS, or to report suspected adverse drug events, please call Virbac at 1-855-647-3747.

P-2001-F-US-1

Lot:

Exp:

NOT APPROVED BY FDA - Legally marketed as an FDA Indexed Product under MIF 900-013.  
FOR USE IN FERRETS ONLY. Extra-label use is prohibited.  
This product is not to be used in animals intended for use as food for humans or other animals.

## Suprelorin® F

(DESLORELIN ACETATE)

4.7 mg IMPLANT



**CONTENTS:** One (1) implant pre-loaded in one (1) sterile implanting needle

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.