



BOOKLET PAGES:




Cover Page 1




Page 2

Page 7




Page 6


Page 3



BOOKLET PAGES:




Cover Page 1




Page 2

Page 7




Page 6

Page 3



Page 4

Page 5



PAGE 4

COVER, PAGE 1

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each 7.5 mL)</b> Loperamide HCl 1 mg</p> <p><b>Purpose</b> Anti-diarrheal</p> <p><b>Use</b> controls symptoms of diarrhea, including Traveler's Diarrhea</p> <p><b>Warnings</b>  <b>Allergy alert:</b> Do not use if you have ever had a rash or other a allergic reaction to loperamide HCl.  <b>Heart alert:</b> Taking more than directed can cause serious heart problems or death.  Do not use if you have bloody or black stool.  <b>Ask a doctor before use if you have:</b>  fever, mucus in the stool, a history of liver disease.  <b>Ask a doctor or pharmacist before use if you are:</b>  taking antibiotics.  When using this product (tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if:</b>  symptoms get worse  diarrhea lasts for more than 2 days  you get abdominal swelling or bulging. These may be signs of a serious condition.</p> <p><b>Directions</b>  drink plenty of clear fluids to help prevent dehydration caused by diarrhea.  Find right dose on chart. If possible, use weight to dose; otherwise use age.  shake well before using.  use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.</p>	<p><b>Drug Facts (continued)</b></p> <p>■ mL = milliliter</p> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td>30 mL after the first loose stool; 15 mL after each subsequent loose stool, but no more than 60 mL in 24 hours</td> </tr> <tr> <td>children 9-11 years (60-95 lbs)</td> <td>15 mL after first loose stool; 7.5 mL after each subsequent loose stool, but no more than 45 mL in 24 hours</td> </tr> <tr> <td>children 6-8 years (40-59 lbs)</td> <td>15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours</td> </tr> <tr> <td>children 2-5 years (34 to 47 lb)</td> <td>ask a doctor</td> </tr> <tr> <td>children under 2 years (up to 33 lb)</td> <td>do not use</td> </tr> </table>	adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool, but no more than 60 mL in 24 hours	children 9-11 years (60-95 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool, but no more than 45 mL in 24 hours	children 6-8 years (40-59 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours	children 2-5 years (34 to 47 lb)	ask a doctor	children under 2 years (up to 33 lb)	do not use	<p><b>Drug Facts (continued)</b></p> <p><b>Other information</b>  ■ each 30 mL contains: sodium 16 mg  store between 20-25°C (68-77°F)  do not use if printed inner or outer neckband is broken or missing.</p> <p><b>Inactive ingredients</b> acetylsalicylic acid, caramel color, D&amp;C Yellow No. 10, FD&amp;C Blue No. 1, flavor, glycerin, microcrystalline cellulose and croscarmellose sodium, propylene glycol, purified water, simethicone emulsion, sodium benzoate, sucralose, titanium dioxide, xanthan gum.</p> <p><b>Questions or comments?</b>  call 1-877-895-3665 (toll-free) or 215-273-8755 (in PA)</p> <p><b>Imodium A-D</b>  Loperamide HCl, Anti-Diarrheal  1 mg per 7.5 mL  www.imodium.com</p>
adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool, but no more than 60 mL in 24 hours												
children 9-11 years (60-95 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool, but no more than 45 mL in 24 hours												
children 6-8 years (40-59 lbs)	15 mL after first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours												
children 2-5 years (34 to 47 lb)	ask a doctor												
children under 2 years (up to 33 lb)	do not use												

PAGE 2

PAGE 3

-----  
**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**  
-----

/s/  
-----

VALERIE S PRATT  
04/12/2017