Adapted from Washington State Department of Health Form DOH 347-102

Fax Page 1 To:

Spokane Regional Health District (509) 324-1468 (Confidential FAX line)



CONFIDENTIAL SEXUALLY TRANSMITTED INFECTION (STI) CASE REPORT
Report STIs within 3 work days (WAC 246-101-101/301)

PATIENT INFORMATION												
LAST NAME FIRST N			IRST NAME	AME			MIDDLE NAN	DAT	DATE OF BIRTH			
									мо		PAY	YR
ADDRESS (Unhoused or	unstably housed i	n the past 3	months)	CITY	,				STATE	ZIP COD	E
TELEPHONE		EMAIL		ENGL	ISH SP	EAKING	G? 🗌 Yes 🔲 N	o *instruct pg.		IAGNOSIS	DATE	
				Prefe	rred La	nguag	e (Code*: L)	N	10	DAY	YR
SEX ASSIGNED	GENDER IDENT		l	HNICITY			RACE CATEGO White	RY (check	all that a	apply)*:		structions on page 3
AT BIRTH			I	Non-Hispanic		☐ Black			☐ Asian ☐ Other			
							☐ American Ir	iska Nat	-			
☐ Intersex	er 🗌 Refused			☐ Refused		☐ Native Haw	ner Pacif	er Pacific Islander Refused				
Refused							EXTENDED RA	CE CODE	S)*: R	R	R	R R
CURRENTLY	REASON FOR E	XAM (check one):	1	OF SEX PAI			call that apply):	HIV STA			AIDS Case Report	CURRENTLY
Yes	PREGNANT? Exposed to I		☐ Male	_ 0					ous positive		ON PrEP?	
□ No	Symptomatic			lale ☐ Transger binary / ☐ Other			ender FTIVI		IIV diagnosis at this visit* ive HIV test at this visit		☐ Yes ☐ No	
☐ Unk. ☐ NA	□ Routine Exai	n (No Symptoms)	_	derqueer 🔲 Unknow			I				□NA	
DIAGNOSIS - D	DISEASE											
GONORRHEA (la	ab confirmed)								SYPHILI	S		
DIAGNOSIS (che	-		•	TREATMENT (check all prescribed):					STAGE (check one):			
☐ Asymptomatic Symptomatic		Cervix	☐ Ceftriaxone: ☐ 250 mg ☐ 500 mg ☐ 1 g				1 g	☐ Primary (Chancre, etc.)☐ Secondary (Rash, etc.)				
	matory Disease			☐ Cefixime: ☐ 400 mg ☐ 800 mg ☐ Azithromycin: ☐ 1 g ☐ 2 g					Early Latent (< 1 year)			
Ophthalmia				☐ Doxycycline: ☐ 100 mg BID x 7 days					Unknown Duration or Late			
☐ Disseminated☐ Other Compli		Pharynx		Gentamicin					☐ Congenital			
		'		☐ Gemifloxacin: ☐ 320 mg ☐ Other:					MANIFESTATIONS (check all that apply):			
Date Tested:		Other:		te Prescrib	 ed:				☐ Neurologic ☐ Otic ☐ Ocular ☐ Tertiary			
CHLAMYDIA (la	b confirmed)	·							TREATN	/IENT (che	ck one):	
DIAGNOSIS (che		SITES (all that a	pply): TR	EATMENT (check	all pres	scribed):			•	.4 MU IM x 1	
☐ Asymptomatic				☐ Azithromycin: ☐ 1 g					☐ 2.4 MU IM x 3			
Symptomatic, Uncomplicated				☐ Doxycycline: ☐ 100 mg BID x 7 days					Doxycycline: ☐ 100 mg BID x 14 days			
☐ Pelvic Inflammatory Disease ☐ Ophthalmia		☐ Urine ☐ Rectum		☐ Levofloxacin: ☐ 500 mg daily x 7 days ☐ Other:				☐ 100 mg BID x 28 days				
☐ Other Compl	ications:	Pharynx		other				Benzathine 50,000 units/kg IM x 1 PCN-G: 50,000 units/kg IM x 3				
		☐ Vagina	į						PCN-G:			_
Date Tested:		☐ Ocular☐ Other:	¦ D	ate Prescrib	ed:				Aqueou Crystalli	+	8-24 MU/day or 10-14 davs	
HERPES SIMPLE	×				R DISE	ASES			Penicilli		л 10-14 days	
DIAGNOSIS		LABORATORY CO	NFIRMATIO		ancroic				Other: _			
_ ` '		☐ Yes		☐ Granuloma Inguinale				Date Prescribed:				
□ Neonatal □ No □ Lymphogranuloma Venereum □ Date Prescribed: □ Lymphogranuloma Venereum □ Lymphogranu												
		treatment by eithe			rson or	hy nre	scribing medicat	ion for na	tionts to	give to the	air say nartna	rs (saa sida 2
for additional info		treatment by eithe	i tieating pa	rtileis ili-pe	13011 01	by pie	F			-	•	-
for additional information). ☐ In-person evaluation - Number of partners treated following medical evaluation: ☐ Turn over for Partner Treatment Plan Instructions												
		Number of partne ir partner(s):					ovided expedited not recommende					
REPORTING CL	LINIC INFORMA	ATION										
DATE	FACILITY NA	AME				DIAGN	IOSING CLINICIA	AN				
ADDRESS				CITY					STATE		ZIP	
PERSON COMPLETING FORM				TELEPHONE			EMA	.IL		1		

PARTNER MANAGEMENT PLAN INSTRUCTIONS

Gonorrhea or Chlamydia Infection: Partner Treatment

All partners should be treated as if they are infected.

The Washington State Department of Health strongly encourages providers to take responsibility to ensure partner treatment for heterosexuals, by examining and treating all patient's sex partners from the previous 60 days.

If an examination is not possible, providers should offer medication for all sex partners whom patients are able to contact.

Spokane Regional Health District may be able to provide free medication to your patient to give their partner(s), if resources permit. Please contact your local health jurisdiction to report cases and inquire about partner management resources, possibly including EPT medications.

Spokane Regional Health District recommends you refer all **MSM patients** and **all patients with syphilis or newly diagnosed HIV** to the health department for help notifying partners to ensure that partners receive medication; the opportunity to test for HIV, syphilis, gonorrhea, and chlamydia; and evaluation for HIV Pre-Exposure Prophylaxis (PrEP). Please inform the patient that the health department will contact them to assist with partner notification.

Although the Health Department requests that you refer patients with these risks to us, we also ask that you make every effort to help patients assure that their partners are treated.

Complete the partner management plan on the Confidential Sexually Transmitted Infection Case Report FAX form to define a partner management plan.

For copies of this case report or questions on how to fill it out, call the Spokane Regional Health District: (509) 324-1494.

Other STIs: Partner Treatment

All patients with infectious syphilis, chancroid, LGV, or granuloma inguinale are routinely contacted by public health staff. Patients diagnosed with genital herpes should be advised to notify their sex partners and should be informed that their partners should contact their provider for testing. Per CDC treatment guidelines, sex partners of patients who are diagnosed with early syphilis (primary, secondary, or early latent) and may be incubating disease should be treated regardless of test results. Alternative treatment for penicillin allergy among non-pregnant patients, such as an appropriate dosage of doxycycline, can also be found in the guidelines.

RECOMMENDED REGIMENS FOR ANTIMICROBIALS LISTED ON REPORTS*

GONORRHEA -- Uncomplicated

Ceftriaxone 500 mg IM as a single dose for persons weighing < 150 kg (330 lbs), or 1 g IM as a single dose for persons weighing > 150 kg (330 lbs) †

Alternatives for uncomplicated infections of the cervix, urethra, or rectum:

Gentamicin 240 mg IM as a single dose PLUS azithromycin 2 g PO as a single dose $\bf OR$ Cefixime 800 mg orally as a single $\bf dose^{\dagger}$

- † If treating with ceftriaxone or cefixime, and chlamydia infection has not been excluded, providers should treat for chlamydia with doxycycline 100 mg PO BID for 7 days. During pregnancy, azithromycin 1 g PO as a single dose is recommended to treat chlamydia.
- [‡] Ceftriaxone is the only recommended treatment for pharyngeal gonorrhea. Consult an infectious disease provider or other STI expert for assistance if alternative treatment is required.

CHLAMYDIA -- Uncomplicated

Doxycycline 100 mg PO BID for 7 days **OR** Azithromycin 1g PO as a single dose

Alternatives:

Erythromycin (base) 500 mg PO QID for 7 days **OR** Ethylsuccinate 800 mg PO QID for 7 days **OR**

Ofloxacin 300 mg PO BID for 7 days **OR**

Levofloxacin 500 mg PO for 7 days

SYPHILIS -- PRIMARY, SECONDARY, OR EARLY LATENT (<1 YEAR)

Benzathine penicillin G 2.4 million units IM in a single dose

SYPHILIS -- LATE OR UNKNOWN DURATION

Benzathine penicillin G 2.4 million units IM for 3 doses at 1 week intervals

DOH 347-102, updated 01/17/2023. For persons with disabilities, this document is available on request in other formats. To submit a request, please call 1-800-525-0127 (TDD/TTY 1-800-833-6388).

^{*} Refer to "STD Diagnostic and Treatment Guidelines" or the Centers for Disease Control and Prevention's (CDC's) website (https://www.cdc.gov/std/treatment/default.htm) for further information on treating pregnant patients, infections of the pharynx, patients with allergies, treatment of infants, and other details.

List of Preferred Languages:

Instructions: Complete the *Preferred Language* section if the patient's preferred language is <u>not</u> English. Patient's preferred language shall be identified by the patient and reported using one of the below categories.

Language Identity Codes:

(L18) Karen	(L35) Sign languages
(L19) Khmer/Cambodian	(L36) Somali
(L20) Kinyarwanda	(L37) Spanish/Castilian
(L21) Korean	(L38) Swahili/Kiswahili
(L22) Kosraean	(L39) Tagalog
(L23) Lao	(L40) Tamil
(L24) Mandarin	(L41) Telugu
(L25) Marshallese	(L42) Thai
(L26) Mixteco	(L43) Tigrinya
(L27) Nepali	(L44) Ukrainian
(L28) Oromo	(L45) Urdu
(L29) Panjabi/Punjabi	(L46) Vietnamese
(L30) Pashto	(L77) Other language
(L31) Portuguese	(L88) Patient declined to respond
(L32) Romanian/Rumanian	(L99) Unknown
	(L19) Khmer/Cambodian (L20) Kinyarwanda (L21) Korean (L22) Kosraean (L23) Lao (L24) Mandarin (L25) Marshallese (L26) Mixteco (L27) Nepali (L28) Oromo (L29) Panjabi/Punjabi (L30) Pashto (L31) Portuguese

(L33) Russian

(L34) Samoan

Race Category and Identity Instructions:

Instructions: The patient's race may either be reported in the general *Race Category* section <u>or</u> the *Extended Race* section on page 1 of the case report. When completing the *Extended Race* section, please report using the codes provided in the list below. If the patient self-identifies as more than one race, each race code shall be reported within the *extended race* section. **Unknown** race can be documented within either one of the race sections.

Extended Race Codes:

(L16) Hmong

(L17) Japanese

(R01) Afghan	(R22) Fijian	(R43) Mexican/Mexican American	(R64) Tongan
(R02) Afro-Caribbean	(R23) Filipino	(R44) Middle Eastern	(R65) Ugandan
(R03) Alaska Native	(R24) First Nations	(R45) Mien	(R66) Ukrainian
(R04) American Indian	(R25) Guamanian or Chamorro	(R46) Moroccan	(R67) Vietnamese
(R05) Arab	(R26) Hmong/Mong	(R47) Native Hawaiian	(R68) White
(R06) Asian	(R27) Indigenous-Latina/o/x	(R48) Nepalese	(R69) Yemeni
(R07) Asian Indian	(R28) Indonesian	(R49) North African	(R77) Other race
(R08) Bamar/Burman/Burmese	(R29) Iranian	(R50) Oromo	(R99) Unknown
(R09) Bangladeshi	(R30) Iraqi	(R51) Pacific Islander	
(R10) Bhutanese	(R31) Japanese	(R52) Pakistani	
(R11) Black or African American	(R32) Jordanian	(R53) Puerto Rican	
(R12) Central American	(R33) Karen	(R54) Romanian/Rumanian	
(R13) Cham	(R34) Kenyan	(R55) Russian	
(R14) Chicana/o or Chicanx	(R35) Khmer/Cambodian	(R56) Samoan	
(R15) Chinese	(R36) Korean	(R57) Saudi Arabian	
(R16) Congolese	(R37) Kuwaiti	(R58) Somali	
(R17) Cuban	(R38) Lao	(R59) South African	
(R18) Dominican	(R39) Lebanese	(R60) South American	
(R19) Egyptian	(R40) Malaysian	(R61) Syrian	
(R20) Eritrean	(R41) Marshallese	(R62) Taiwanese	
(R21) Ethiopian	(R42) Mestizo	(R63) Thai	